



Trinity Road, Sutton Coldfield
West Midlands B75 6TJ

www.coppice.bham.sch.uk

Tel No: 0121 4647022

Fax No: 0121 4642180

e-mail: enquiry@coppice.bham.sch.uk

PLEASE CHECK YOUR CHILD'S HEAD VERY CAREFULLY AND READ THE FOLLOWING INFORMATION.

WHAT ARE HEADLICE?

Head lice are small (smaller than a match head) insects that can live only on the head.

WHO CAN GET HEAD LICE?

Anyone can get head lice – they are not a sign of being dirty. Head lice occur wherever there are children at home or in school. Hair can be long or short.

WHAT ARE THE SYMPTOMS OF HEAD LICE AND WHEN DO THEY APPEAR?

Symptoms are itching and scratching of the scalp usually 2-3 weeks after head lice get hold of the head.

HOW ARE HEAD LICE SPREAD?

Head lice are passed by direct contact of heads, or when personal items (combs, brushes, hats, scarves, headphones, jackets and clothing) and bedding are shared. Head lice only crawl and cannot jump from person to person. Head lice do not spread to pets and you cannot get them from pets.

HOW DO HEAD LICE LIVE?

Head lice lay eggs close to the scalp. The eggs are firmly stuck to the hair, and difficult to see with the naked eye. Eggs can only hatch on the head itself, not elsewhere. Eggs take 7-10 days to hatch and a further 7-14 days to become fully grown, and start laying eggs. Empty egg sacs are white and shiny, and may be found further along the hair, as the hair grows out. Head lice feed by sucking blood through the scalp.

HOW ARE HEAD LICE DETECTED?

Shampoo hair and apply hair conditioner liberally to make hair slippery. Wet hair makes it difficult for the lice to keep a grip on the hair. Head lice are then easily removed. Comb hair from the scalp outwards with a fine comb (or a 'nit' comb available from chemists) for ten minutes over a sink or disposable towel. Lice are present if they fall out or stick to the comb.

Make sure the teeth of the comb slot into the hair at the roots with every stroke. The comb should be washed with water or wiped clean with a paper napkin after each stroke. If no head lice are detected then repeat this procedure each week when the hair is washed, as prevention.

HOW ARE HEAD LICE TREATED?

'Wet combing method': This method does not use chemicals and is the same as for detecting head lice. Just repeat the method for detecting head lice every day for the first week, then every third day for the second week and once a week for the third and fourth weeks.

The 'Wet' combing method to detect and remove head lice is effective and recommended.

N.B Medicated lotions should only be used as a last resort.

HOW CAN THE FURTHER SPREAD OF HEAD LICE BE AVOIDED?

All family members with head lice should be treated promptly.

Recently worn clothing (including hats, scarves, jackets and pyjamas) should be washed. Non-washable items such as sports headgear, headphones, stuffed animals and pillows should be stored in plastic bags for a week and then used again.

HOW CAN THE SPREAD OF HEAD LICE BE PREVENTED?

Comb hair, at least once a day, with a fine tooth comb. Brushing alone is not effective in avoiding head lice.

WHERE DO I GET FURTHER ADVICE?

The school nurse or your own doctor will provide you with additional advice.

School Nurses
Falcon Lodge Clinic
Churchill Road
Sutton Coldfield
West Midlands
B76 7LB

Tel: 0121 465 5562