



# Intraabdominal Hemorrhage Due to Hepatic Cyst Rupture During CPR: Case Report

## Kardiyopulmoner Canlandırma Sırasında Hepatik Kist Rüptürüne Bağlı İntraabdominal Hemoraji: Olgu Sunumu

Kardiyopulmoner Canlandırma Sırasında İntraabdominal Hemoraji / Intraabdominal Hemorrhage During CPR

Mehmet Tokaç<sup>1</sup>, Gözde Bumin Aydın<sup>2</sup>, Fahri Yetişir<sup>1</sup>, Murat Baki Yıldırım<sup>1</sup>, Mehmet Kılıç<sup>3</sup>  
<sup>1</sup>Ankara Atatürk Eğitim Araştırma Hastanesi Genel Cerrahi Kliniği,  
<sup>2</sup>Dışkapı Yıldırım Beyazıt Eğitim Araştırma Hastanesi Anestezi Kliniği,  
<sup>3</sup>Yıldırım Beyazıt Üniversitesi Genel Cerrahi ABD, Ankara, Türkiye

### Özet

Kardiyopulmoner canlandırma, seçilmiş hasta popülasyonlarında sağ kalımı arttırmaktan sorumludur. Ancak bu hayatta kalan grup içinde external göğüs kompresyonuna bağlı postresuscitative beklenmedik olaylar olabilir. Visceral organ yaralanmaları bunların bir kısmını oluşturur. Karaciğer kistleri genellikle otozomal dominant polikistik böbrek hastalığına eşlik eder. Kronik renal yetmezliği olan bu hastalarda, üremiye bağlı kanamaya yatkınlık veya diyaliz sırasında kullanılan antikoagulan tedavinin etkisi olarak visceral organ yaralanması sonucu kanama daha kolay olabilir. Böyle hastalarda CPR sonucu devam eden ısrarlı hemodinamik instabilite multidisipliner olarak değerlendirilmelidir. Bu yazımızda, kardiyopulmoner canlandırma sonrası karaciğer kist rüptürüne bağlı masif intraabdominal hemoraji gelişen periton diyaliz hastası sunuldu.

### Anahtar Kelimeler

Kardiyopulmoner Canlandırma (CPR); Karaciğer Yaralanması; Polikistik Karaciğer Hastalığı; Polikistik Böbrek Hastalığı

### Abstract

Cardiopulmonary Resuscitation (CPR) improves survival in selected patient populations. However, in the surviving group, postresuscitative unexpected complications can occur due to external chest compression. Visceral organ injuries are a part of their. Liver cysts are usually accompanied by an autosomal dominant polycystic kidney disease. In patients with chronic renal failure, bleeding tendency in uremia is due. As the effect of anticoagulant drugs used during dialysis. Also, visceral organ bleeding due to injury may be easier. In such patients, ongoing hemodynamic instability after CPR, should be regarded as multidisciplinary. In this article, massive intra-abdominal hemorrhage due to rupture of hepatic cyst after CPR, patient with peritoneal dialysis treatment is presented.

### Keywords

Cardiopulmonary Resuscitation (CPR); Liver Injury; Polycystic Hepatic Disease; Polycystic Kidney Disease

DOI: 10.4328/JCAM.1187

Received: 23.07.2012 Accepted: 12.09.2012 Printed: 01.03.2014

J Clin Anal Med 2014;5(2): 122-4

Corresponding Author: Mehmet Tokaç, Subayevleri Mah. Güzin Sok. 36/2 Ankara, Türkiye.

T.: +905052914623 E-Mail: drtokac@gmail.com

## Introduction

Polycystic hepatic disease (PHD) is a rare clinical disorder. Its prevalence is 0.05-0.13% in autopsies. [1]. It is usually together with autosomal dominant polycystic kidney disease (ADPKD) and it is the most common extra renal symptom of polycystic hepatic disease [2,3]. Liver cyst incidence rises with age and has an incidence of 60-75% in the end stage renal failure patients. Hepatic cysts are usually asymptomatic and severe complications like hemorrhage and infection which are rarely seen [4]. External chest compressions performed during cardiopulmonary Resuscitation (CPR) could cause different complications. Sternum and costal traumas takes the first place [5]. But the trauma in intraabdominal visceral organs associated with CPR is a rare complication with limited literature [6;7].

In our case we report a successful CPR in a patient with polycystic kidney disease causing intraabdominal hemorrhage due to hepatic cyst rupture.

## Case Report

The 40 year old patient has had peritoneal dialysis for five years. After the peritoneal dialysis he had a pain in epigastric at left sternal region. He fainted and referred to a government hospital emergency department. He was accepted to intensive care unit (ICU) as he was thought to have a heart attack. In ICU as he had an arrest, he required CPR and his trachea was intubated. After 24 hours he was referred to our hospital for hemodialysis and for clinical follow up with a helicopter ambulance. He was admitted to Intensive Care Unit and was connected to mechanical ventilation using CPAP (continuous pressure assisted ventilation) mode. Central venous catheter and arterial line was induced to the patient. His blood pressure was 130/70 mmHg, his heart rate was 109beats/min, and his temperature was 36.7°C, with a breathing rate of 16/min and 5 cm H<sub>2</sub>O of central venous pressure (CVP). Relevant laboratory investigation included as it is below indicated: hemoglobin level 16.7g/dl, platelet count 209X10<sup>9</sup>/l, white cell count 18.3X10<sup>9</sup>/l with normal differential count. Serum urea amounted to 142 mg/dl, creatinine 13.48 mg/dl, sodium 134 mmol/l and potassium 5.6 mmol/l. Liver biochemistry tests were abnormal: ALT 285 U/l, AST 169 U/l. Blood gas analysis revealed; pH 7.30, pCO<sub>2</sub> 46 mmHg, base excess -5.3 mmol/l, bicarbonate 21.6 mmol/l, pO<sub>2</sub> 135 mmHg, O<sub>2</sub> saturation 99%. International normalized ratio (INR) was so high that it could not be measured. Troponin I was 50ng/dl (0.02-0.06). He was replaced with 4 units of fresh frozen plasma. His control INR was 1.5. His peritoneal fluid was discharged with peritoneal dialysis catheter and the peritoneal fluid seemed clear. He was scheduled to daily hemodialysis programme.

Low molecular weight heparin 5000 iu was given to the patient twice a day. After stable vital signs for 24 hours and satisfactory breathing, he was extubated. After 40 hours of admission to the ICU his blood pressure dropped and tachycardia began. Blood pressure was 95/50 mmHg and heart rate was 123/min. The CVP was 2 cmH<sub>2</sub>O. His hemoglobin level was 11.6gr/dl and hematocrite was 31.8%, INR was 1.35. He did not have abdominal pain and did not have abdominal rebound, defence or tenderness. Nasogastric irrigation was performed to rule out gastric bleeding. hemorrhage was not detected. His rectal digital

examination was also normal. As his hemoglobin level dropped to 7.5 gr/dl, he had an abdominal computerized tomography (CT). In his abdominal CT a 19 mm perihepatic and 15 mm perisplenic, 15 mm right paracolic and 10 mm left paracolic fluid was detected. In abdominal CT liver cysts 45X23 mm in left lobe and 30X14 mm in right lobe were detected. [Fig1] Both of his kidneys were also in polycystic shape. Again peritoneal dialysis catheter was irrigated and hemoragic fluid was detected referring to the intraabdominal hemorrhage. The patient was operated under general anesthesia with the presumptive diagnosis of intraabdominal hemorrhage. There was a 1500cc of hemoragic fluid between splenic and subhepatic space. During abdominal exploration we saw that the liver was multicystic. The cyst at the segment 3 in the inferior part of the liver was ruptured and actively bleeding. We used primary sutures to stop the bleeding but the liver was so fragile because of cystic structure thus to control bleeding omentopexia was performed. [Fig 2] Four units of erythrocyte suspension was infused to the patient during the operation. After the operation the patient was followed up in ICU. He did not have any bleeding postoperatively and his hemoglobin levels was stable. We extubated the patient at his first ICU day. Postoperatively hemodialysis programme continued. Three days later after the operation, we added colpodogrel 75 mg to his medical order. Postoperative 5th day coronary angiogram was performed and 100% occlu-



Figure 1. The axial CT scan shows free fluid and polycystic liver and kidney disease.




Figure 2. View of ruptured cyst in left liver lobe, that was sutured in order to establish hemostasis.

sion was found at his right coronary artery and was stented. He was discharged from ICU on the eight day of operation. He was discharged from hospital at 11th day postoperatively with a 3 day/week dialysis programme.

## Discussion

Hepatic cysts are the most common extrarenal manifestation of ADPKD. In ADPKD hepatic cysts prevalence rises with age. It has 10% prevalence in 20-29 age interval, but it has a prevalence of 75% in patients who are older than 60 years [6]. Because of this, hepatic cysts can be detected in dialysis patients more commonly [7]. Rarely cyst infection [8], hemorrhage and intraabdominal hemorrhage due to hepatic cyst rupture [9], torsion of the cyst [5], portal hypertension [10], hepatitis due to biliary tract compression [11] are the complications.

Skeletal complications due to CPR usually don't cause any mortality but rarely visceral trauma could threaten life of the patient [12]. Intraabdominal complications about Resuscitation can be due to trauma of hepatic, splenic or other gastrointestinal organs. The most common visceral trauma in literature is hepatic trauma [12]. A study by Kriccher et al about extrahospital CPR complications determined in 705 of 2187 autopsies had a hepatic rupture incidence of 2.1% (n=15). [5] Another single centered study showed 0.6% hepatic rupture in patients resuscitated for cardiac arrest. Among 15 hepatic ruptures, 6 were detected in autopsies. In both of the studies there is no information about the hepatic pathologies of the patients. In the current literature there is no peritoneal dialysis patient developing intraabdominal hemorrhage because of hepatic cyst rupture after CPR.

Hepatic ruptures developing after CPR could be seen in left lobe because of the neighbourhood of the heart to the left lobe of the liver. Right lobe ruptures are also seen in the literature [13]. There are some case reports about coagulability or thrombotic treatments causing hemorrhagic complications [14]. Risk factor for bleeding in chronic hemodialysis patients is uremic bleeding tendency, repeated systemic anticoagulation during dialysis procedures [14]. There may be some difficulties in diagnosis with hepatic trauma in sedated patients. Hypovolemia secondary to bleeding caused by trauma could rise slowly and can be similar to volume depletion caused by ischemia reperfusion due to CPR. Hemodynamic instability after Resuscitation could cause misdiagnosis. Hematocrit level is a quantitative measurement.

Contrasted computerised tomography (CT) is a useful diagnostic method in intraperitoneal hemorrhage [13]. In our case uncontrasted CT was taken in emergent conditions because of patient's uremia and we could only see intraabdominal fluid. Intraabdominal fluid could be normally seen in patients with peritoneal dialysis. Our diagnosis was made by aspiration of hemorrhagic fluid from his peritoneal dialysis catheter.

The goal of the treatment of hepatic parenchymal traumas is to control the bleeding, debridement of devitalized hepatic tissue and the drainage of perihepatic area. Omentoplasty is a functional method to achieve hemostasis when primary suturation is not enough for controlling bleeding [15].

In conclusion Emergent and persistent chest compressions are vital during CPR but they can cause hepatic trauma. Multiple

liver cysts are tend to cause bleeding due to hepatic rupture. Patients with ADPKD with uremic coagulability could potentiate this problem. We should be alert to patients during ICU follow up with hemodynamic instability and hematocrit impairment after CPR and radiological imaging techniques should be administered to the patient. Timing of diagnosis and appropriate treatment are life saving factors in hepatic traumas.

## Competing interests

The authors declare that they have no competing interests.

## References

1. Karhunen PJ, Tenhu M. Adult polycystic liver and kidney diseases are separate entities. *Clin Genet* 1986;30(1):29-37.
2. Bistriz L, Tamboli C, Bigam D, Vincent G. Polycystic Liver Disease: Experience at a Teaching Hospital. *Am J Gastroenterol* 2005;100(10):2212-7.
3. Krischer JP, Fine EG, Davis JH, Nagel EL. Complications of cardiac resuscitation. *Chest* 1987;92(2):287-91.
4. Vock R. Liver and splenic ruptures as a complication of heart pressure massage. *Beitr Gerichtl Med* 1992;50:193-203.
5. Smekal D, Johansson J, Huzevka T, Rubertsson S. No difference in autopsy detected injuries in cardiac arrest patients treated with manual chest compressions compared with mechanical compressions with the LUCASTM device-A pilot study. *Resuscitation* 2009;80(10):1104-7.
6. Milutinovic J, Fialkow PJ, Rudd TG, Agodoa LY, Phillips LA, Bryant JL. Liver cysts in patients with autosomal dominant polycystic kidney disease. *Am J Med* 1980;68(5):741-4.
7. Carels RA, Bommel EFH. Ruptured giant liver cyst: a rare cause of acute abdomen in a haemodialysis patient with autosomal dominant polycystic kidney disease. *Neth J Med* 2002;60(9):363-5.
8. Srinivasan R. Polycystic liver disease: An unusual cause of bleeding varices. *Dig Dis Sci* 1999;44(2):389-92.
9. Cappell MS. Obstructive jaundice from benign, nonparasitic hepatic cysts: Identification of risk factors and percutaneous aspiration for diagnosis and treatment. *Am J Gastroenterol* 1988;83(10):93-6.
10. Hoke RS, Chamberlain D. Skeletal chest injuries secondary to cardiopulmonary resuscitation. *Resuscitation* 2004;63(3):327-38.
11. Kouzu H, Hase M, Kokubu N, Nishida J, Kawamukai M, Usami Y, et al. Delayed visceral bleeding from liver injury after cardiopulmonary resuscitation. *J Emerg Med* 2012;43(4):245-8.
12. Chung TK, Chen KS, Yen CL, Chen HY, Cherng WJ, Fang KM. Acute abdomen in a haemodialysed patient with polycystic kidney disease--rupture of a massive liver cyst. *Nephrol Dial Transplant* 1998;13(7):1840-2.
13. Fehmi C, Balık A, Polat Y, Yıldırım I, Boyuk A, Oren D. Hepatic injuries; our surgical treatment experience. *Ulus Travma Acil Cerrahi Derg* 2001;7(3):185-8.