

Drugs to Avoid with Myasthenia Gravis

Antibiotics	Heart medications	Anesthesia	Brain/Nerve	Others
ampicillin	Quinidine	Procainamide	Lithium	Timolol eye drops
Amoxicillin	Quinine	Succinylcholine	Phenytoin	Cortisones
Penicillin	Procainamide	Curare derivatives	Gabapentin	Penicillamine
Imipenem	Statins		Botox	Iodinated contrast
Aztreonam	Atenolol		Nicotine	Magnesium
Ofloxacin	Metoprolol		Methocarbamol	Interferon alpha
Levofloxacin	Sotalol			
Ciprofloxacin	Propranolol			
Erythromycin	Pindolol			
Clindamycin	Nebivolol			
Azithromycin	Nadolol			
Clarithromycin	Labetalol			
Amikacin	Esmolol			
Gentamycin	Carvedilol			
Tobramycin	Bisoprolol			
Kanamycin	Acebutolol			
Neomycin	Amlodipine			
Streptomycin	Verapamil			
	Diltiazem			
	Nifedipine			
	Felodipine			

Red: Strong evidence of harmful effect

Yellow: Few case reports of harmful effect

Blue: Should be avoided in spite of weak clinical evidence

NB: to make it a patient friendly list, names of all common individual medications were listed instead of pharmaceutical group listing so you can copy this page and give it to the patient as a reference.