

# 2017 Virginia Prevention Needs Assessment Survey



Survey Results for:  
Chesterfield County

Sponsored by:  
SAFE, Inc.  
P.O. Box 40  
Chesterfield, Virginia 23832  
804-796-7100

## 2017 Prevention Needs Assessment Survey Profile Report

This report summarizes the findings from the Prevention Needs Assessment (PNA) Survey that was conducted during 2017. The results are presented along with comparisons to national data sources such as the Monitoring the Future Survey (only grades 8, 10, and 12 are surveyed) and the Bach Harrison Norm (BH Norm), which consists of a large, weighted, nationwide sample.

The survey was designed to assess students' involvement in a specific set of problem behaviors, as well as their exposure to a set of scientifically validated risk and protective factors. The risk and protective factors have been shown to influence the likelihood of academic success, school dropout, substance abuse, violence, and delinquency among youth.

Table 1 contains the characteristics of the students who completed the survey from your community. When using the information in this report, please pay attention to the number and

### Contents:

- Introduction
- The Risk and Protective Factor Model of Substance Abuse Prevention
- Building a Strategic Prevention Framework
- Validity Measures
- How to Read the Charts
- Tools for Assessment and Planning
- ATOD and Antisocial Behavior Charts
- Risk and Protective Factor Charts
- Risk and Protective Factor Scale Definitions
- Data Tables
- Drug Free Communities and Youth Perception of Substance Use Report
- Contacts for Prevention

percentage of students who participated from your community. The sample size for this survey administration was 4,731 students. If 60% or more of the students sample participated, the report is a good indicator of the levels of substance use, risk, protection, and antisocial behavior. If fewer than 60% participated, a review of who participated should be completed prior to generalizing the results to the entire community.

## The Risk and Protective Factor Model of Substance Abuse Prevention

Many states and local agencies have adopted the Risk and Protective Factor Model to guide their prevention efforts. The Risk and Protective Factor Model of Prevention is based on the simple premise that to prevent a problem from happening, we need to identify the factors that increase the risk of that problem developing and then find ways to reduce the risks. Just as medical researchers have found risk factors for heart disease such as diets high in fat, lack of exercise, and smoking; a team of researchers at the University of Washington have defined a set of risk factors for youth problem behaviors.

Risk factors are characteristics of school, community, and family environments, as well as characteristics of students and their peer groups that are known to predict increased likelihood of drug use, delinquency, school dropout, teen pregnancy, and violent behavior among youth. Dr. J. David Hawkins, Dr. Richard F. Catalano, and their colleagues at the University of Washington, Social Development Research Group have investigated the relationship between risk and protective factors and youth problem behavior. For example, they

**Table 1. Characteristics of Participants**

Student Totals						
Total Students	Chesterfield County					
	2014		2016		2017	
	Number	Percent	Number	Percent	Number	Percent
	3745	100	3514	100	4141	100
Grade						
8	1373	39.1	1096	31.2	1940	46.8
10	1161	33.0	1266	36.0	1254	30.3
12	1211	34.5	1152	32.8	947	22.9
Gender						
Male	1730	46.5	1751	50.0	2070	50.2
Female	1990	53.5	1748	50.0	2052	49.8
Ethnicity						
Native American	27	0.8	18	0.5	28	0.7
Asian	135	3.9	149	4.3	167	4.0
African American	876	25.1	833	23.9	898	21.8
Pacific Islander	21	0.6	9	0.3	10	0.2
Hispanic	263	7.5	329	9.4	347	8.4
White	1980	56.7	1707	48.9	2125	51.5
Multi-racial	423	12.1	444	12.7	552	13.4

Table 1. represents the total survey population. Students were given the option to skip questions, and not all students completed the survey. The percentages in remaining tables/figures of this report reflect the percent of students responding to each question, rather than the percent of the total survey population.

# Risk and Protective Factors

## The Risk and Protective Factor Model of Substance Abuse Prevention (Continued)

have found that children who live in families with high levels of conflict are more likely to become involved in problem behaviors such as delinquency and drug use than children who live in families with low levels of family conflict.

Protective factors exert a positive influence or buffer against the negative influence of risk, thus reducing the likelihood that adolescents will engage in problem behaviors. Protective factors identified through research reviewed by Drs. Hawkins and Catalano include social bonding to family, school, community and peers; healthy beliefs and clear standards for behavior; and individual characteristics. For bonding to serve as a protective influence, it must occur through involvement with peers and adults who communicate healthy values and set clear standards for behavior. Research on risk and protective factors has important implications for prevention efforts.

The premise of this approach is that in order to promote positive youth development and prevent problem behaviors, it is necessary to address those factors that predict the problem.

By measuring risk and protective factors in a population, prevention programs can be implemented that will reduce the elevated risk factors and increase the protective factors. For example, if academic failure is identified as an elevated risk factor in a community, then mentoring, tutoring, and increased opportunities and rewards for classroom participation can be provided to improve academic performance. The chart to the right shows the links between the 20 risk factors and the six problem behaviors. The check marks have been placed in the chart to indicate where at least two well designed, published research studies have shown a link between the risk factor and the problem behavior.

Risk Factors for Adolescent Problem Behavior	Problem Behaviors					
	Substance Abuse	Delinquency	Teen Pregnancy	School Drop-Out	Violence	Depression & Anxiety
<b>Community</b>						
Community Laws and Norms Favorable Toward Drug Use, Firearms and Crime	✓	✓			✓	
Perceived Availability of Drugs	✓	✓				
<b>Family</b>						
Poor Family Management	✓	✓	✓	✓	✓	✓
Family Conflict	✓	✓	✓	✓	✓	✓
Sibling Drug Use	✓	✓	✓	✓	✓	✓
Exposure to Adult Antisocial Behavior	✓	✓	✓	✓	✓	✓
Parent Attitudes Favor Antisocial Behavior	✓	✓			✓	✓
Parent Attitudes Favor Drug Use	✓	✓			✓	✓
<b>School</b>						
Academic Failure	✓	✓	✓	✓	✓	✓
Low Commitment to School	✓	✓	✓	✓	✓	✓
<b>Peer / Individual</b>						
Early Initiation of Antisocial Behavior	✓	✓	✓	✓	✓	✓
Early Initiation of Drug Use	✓	✓	✓	✓	✓	✓
Attitudes Favorable to Antisocial Behavior	✓	✓	✓	✓	✓	
Attitudes Favorable to Drug Use	✓	✓	✓	✓	✓	
Perceived Risk of Drug Use	✓	✓	✓	✓	✓	
Interaction with Antisocial Peers	✓	✓	✓	✓	✓	✓
Friend's Use of Drugs	✓	✓	✓	✓	✓	✓
Rewards for Antisocial Behavior	✓	✓	✓	✓	✓	✓
Depressive Symptoms	✓			✓		✓
Gang Involvement	✓	✓			✓	✓



# Building a Strategic Prevention Framework

The survey is an important data source for the Substance Abuse and Mental Health Services Administration (SAMHSA) Center for Substance Abuse Prevention (CSAP) Strategic Prevention Framework (SPF). CSAP created the SPF model to guide states and communities in creating planned, data-driven, effective, and sustainable prevention programs. Each part represents an interdependent element of the ongoing process of prevention coordination.

**Assessment: Profile Population Needs, Resources, and Readiness to Address the Problems and Gaps in Service Delivery.** The SPF begins with an assessment of the needs in the community that is based on data. One of the primary sources of needs assessment data is this Prevention Needs Assessment Survey (PNA). While planning prevention services, communities are urged to collect and use multiple data sources, including archival and social indicators, assessment of existing resources, key informant interviews, and community readiness. The PNA results presented in this Profile Report will help you to identify needs for prevention services. PNA data include adolescent substance use, anti-social behavior, and many of the risk and protective factors that predict adolescent problem behaviors.

**Capacity: Mobilize and/or Build Capacity to Address Needs.** Engagement of key stakeholders at the State and community levels is critical to plan and implement successful prevention activities that will be sustained over time. Some of the key tasks to mobilize the state and communities are to work with leaders and stakeholders to build coalitions, provide training, leverage resources, and help sustain prevention activities.

**Planning: Develop a Comprehensive Strategic Plan.** States and communities should develop a strategic plan that articulates not only a vision for the prevention activities, but also strategies for organizing and implementing prevention efforts. The strategic plan should be based on the assessments conducted during Step 1. The Plan should address the priority needs, build on identified resources/strengths, set measurable objectives, and identify how progress will be monitored. Plans should be adjusted with ongoing needs assessment and monitoring activities.

**Implementation: Implement Evidence-based Prevention Programs and Infrastructure Development Activities.** By measuring and identifying the risk factors and other causal factors that contribute to the targeted problems specified in your strategic plan, programs can be implemented that will reduce the prioritized substance abuse problems. After completing Steps 1, 2, and 3, communities will be able to choose prevention strategies that



## Building a Strategic Prevention Framework (cont'd)

have been shown to be effective, are appropriate for the population served, can be implemented with fidelity, are culturally appropriate, and can be sustained over time. SAMHSA developed an internet tool located at [www.samhsa.gov/nrepp](http://www.samhsa.gov/nrepp) for identifying Best Practice Programs. Another way to access the resources is from this address [www.nrepp.samhsa.gov](http://www.nrepp.samhsa.gov).

**Evaluation: Monitor Process, Evaluate Effectiveness, Sustain Effective Programs/Activities, and Improve or Replace Those That Fail.** Finally, ongoing monitoring and evaluation are essential to determine if the desired outcomes are achieved, assess service delivery quality, identify successes, encourage needed improvement, and promote sustainability of effective policies, programs, and practices. The OPNA allows communities to monitor levels of ATOD use, antisocial behavior, risk, and protection.

**Sustainability and Cultural Competence: Incorporate principles of cultural competence and sustainability in each of the five elements.** At the center of the SPF model, sustainability and cultural competence play a key role in assessment, capacity appraisal, planning, implementation and evaluation, ensuring successful, long lasting prevention programs.

**Sustainability is accomplished by utilizing a comprehensive approach.** States and communities should plan adaptive, flexible programs around a variety of resources, funding, and organizations. An inclusive design helps build sustainable programs and achieve sustainable outcomes. A strategic plan that dynamically responds to changing issues, data, priorities, and resources is more likely to achieve long term results.

Sharing information gathered during the evaluation stage with key stakeholders, forging partnerships and encouraging creative collaboration all enhance sustainability.

**Cultural Competence recognizes unique needs, styles, values and beliefs of the recipients of prevention efforts.** Culturally competent prevention strategies use interventions, evaluations and communication strategies appropriate to their intended community. Cultural issues reflect a range of influences and are not just a matter of ethnic or racial identity. Learning to communicate with audiences from diverse geographic, cultural, economic, social, and linguistic backgrounds can increase program efficacy and ensure sustainable results.

Whether enlisting extended family networks as a prevention resource for single parent households, or ensuring there are resources available to bridge language gaps, cultural competency will help you recognize differences in prevention needs and tailor prevention approaches accordingly.

A one-size-fits-all program is less effective than a program that draws on community-based values, traditions, and customs and works with knowledgeable people from the community to develop focused interventions, communication, and support.

## Validity Measures

**Honesty:** Because the survey was anonymous, and because confidentiality was stressed through the survey's administration process, most of the reasons for students to exaggerate or deny behaviors were eliminated. However, Bach Harrison has built several checks into the data analysis to minimize the impact of students who were either not truthful in their responses or who did not take the survey seriously. Surveys were eliminated from the final data reported in this report for meeting one or more the following five pre-determined dishonesty indicators:

1. In response to a question about whether or not they had been honest in completing the survey, the students indicated that they were "Not Honest At All" in completing the survey.
2. The students indicated that they had used a non-existent, fictitious drug in their lifetime or in the past 30 days.
3. The students reported an impossibly high level of multiple drug use (having used substances on 120 or more occasions in the past 30 days).
4. The students indicated past-month use rates that were higher than lifetime use rates. The student can make one mistake, which is then recoded so that the lifetime is equal to the 30 day value.
5. The students reported an age that was inconsistent with their grade or their school; for example, a 10-year-old 12th grader or 19-year-old 6th grader.

Additionally, if a student did not answer enough of the validity questions to determine whether or not they were honest in their responses, their survey data were also removed from the final analysis presented in this report.

## How to Read the Charts

There are four types of charts presented in this report:

1. Substance use charts
2. Antisocial behavior (ASB) and Gambling charts
3. Risk factor charts
4. Protective factor charts.

Data from the charts are also presented in Tables 3 through 10. Additional data found in later tables are explained at the end of this section.

### Understanding the Format of the Charts

There are several graphical elements common to all the charts. Understanding the format of the charts and what these elements represent is essential in interpreting the results of the PNA survey.

**The Bars** on substance use and antisocial behavior charts represent the percentage of students in that grade who reported a given behavior. The bars on the risk and protective factor charts represent the percentage of students whose answers reflect significant risk or protection in that category. Each set of differently colored bars represents one of the past administrations of the PNA. By looking at the percentages over time, it is possible to identify trends in substance use and antisocial behavior. By studying the percentage of youth at risk and with protection over time, it is possible to determine whether the percentage of students at risk or with protection is increasing, decreasing, or staying the same. This information is important when deciding which risk and protective factors warrant attention.

**Dots and Diamonds** provide points of comparison to larger samples. The dots on the charts represent the percentage of all of the youth surveyed who reported substance use, problem behavior, elevated risk, or elevated protection. Please note that the dot represents the aggregate results of all participating students rather than a random sample of students. The survey results provide considerable information for communities to use in planning

## How to Read the Charts (cont'd)

prevention services. The diamonds represent national data from either the Monitoring the Future (MTF) Survey or the Bach Harrison Norm (BH Norm). The BH Norm was developed by Bach Harrison L.L.C. to provide states and communities with the ability to compare their results on risk, protection, and antisocial measures with more national measures. Survey participants from eight statewide surveys and five large regional surveys across the nation were combined into a database of approximately 460,000 students. The results were weighted to make the contribution of each state and region proportional to its share of the national population. Bach Harrison analysts then calculated rates for antisocial behavior and for students at risk and with protection. The results appear on the charts as BH Norm. In order to keep the BH Norm relevant, it is updated approximately every two years as new data become available.

A comparison to state-wide and national results provides additional information for your community in determining the relative importance of levels of alcohol, tobacco and other drug (ATOD) use, antisocial behavior, risk, and protection. Information about other students in the state and the nation can be helpful in determining the seriousness of a given level of problem behavior. Scanning across the charts, it is important to observe the factors that differ the most from the BH Norm. This is the first step in identifying the levels of risk and protection that are higher or lower than those in other communities. The risk factors that are higher than the BH Norm and the protective factors are lower than the BH Norm are probably the factors that you should consider addressing when planning prevention programs.

### Cut-Points

Before the percentage of youth at risk on a given scale could be calculated, a scale value or cut-point needed to be determined that would separate the at-risk group from the not at-risk group. The Prevention Needs Assessment (PNA) survey was designed to assess adolescent substance use, anti-social behavior, and the risk and protective factors that predict these adolescent problem behaviors. Since the PNA survey has recently been given to over 460,000 youth nationwide, it was possible to select two groups of youth, one that was more at risk for problem behaviors and another group that was less at risk. A cut-point score was then determined for each risk and protective factor scale that best divided the youth

from the two groups into their appropriate group, more at-risk or less at-risk. The criteria for separating youth into the more at-risk and the less at-risk groups included academic grades (the more at-risk group received “D” and “F” grades, the less at-risk group received “A” and “B” grades), ATOD use (the more at-risk group had more regular use, the less at-risk group had no drug use and use of alcohol or tobacco on only a few occasions), and antisocial behavior (the more at-risk group had two or more serious delinquent acts in the past year, the less at-risk group had no serious delinquent acts).

The cut-points that were determined by analyzing the results of the more at-risk and less at-risk groups will remain constant and will be used to produce the profiles for future surveys.

Since the cut-points for each scale will remain fixed, the percentage of youth above the cut-point on a scale (at-risk) will provide a method for evaluating the progress of prevention programs over time. For example, if the percentage of youth at risk for family conflict in a community prior to implementing a community-wide family/parenting program was 60% and then decreased to 50% one year after the program was implemented, the program would be viewed as helping to reduce family conflict.

### Lifetime, 30 Day & Heavy ATOD Use Charts

There are three types of use measured on the ATOD charts.

**Ever-used** is a measure of the percentage of students who tried the particular substance at least once in their lifetime and is used to show the percentage of students who have had experience with a particular substance.

**30-day use** is a measure of the percentage of students who used the substance at least once in the 30 days prior to taking the survey and is a more sensitive indicator of the level of current use of the substance.

**Heavy use** is measured in two ways: *binge drinking* (five or more drinks in a row over the last two weeks), and use of *one-half a pack or more of cigarettes per day*.

### Supplement 1. ATOD Questions

This contains the full wording for the ATOD questions.

## How to Read the Charts (cont'd)

### Antisocial Behavior, Driving and Alcohol, and Gambling Charts

**Antisocial behavior (ASB)** is a measure of the percentage of students who report any involvement during the past year with the two antisocial behaviors listed in the charts.

**Driving and Alcohol** is a measure of the percentage of students who report drinking and driving, or being a passenger in a car where the driver had been drinking in the past 30 days.

### Risk and Protective Factor Charts

Risk and protective factor scales measure specific aspects of a youth's life experience that predict whether he/she will engage in problem behaviors. The scales, defined in Table 2, are grouped into four domains: community, family, school, and peer/individual. The risk and protective factor charts show the percentage of students at risk and with protection for each of the scales. Along with the scales, there are bars that show the percentage of High Risk Youth and percentage of High Protection Youth. High Risk Youth is defined as the percentage of students who have more than a specified number of risk factors operating in their lives. The number of factors is listed on the charts and tables. High Protection Youth is defined as the percentage of students who have more than a specified number of protective factors operating in their lives and is also listed on the tables and graphs.

## Additional Tables

### Additional Tables in this Report

Table 10, Sources and Places of Student Alcohol Use, presents the percentages of how and where students obtained and used alcohol during the past year. The data focus on a subgroup of students who indicated at least one means of obtaining or using alcohol. (Students reporting no alcohol use are not represented.) It is important to note that the table represents a subgroup of users and not the entire survey population. Additionally, the smaller the sample, the more dramatic the influence of a student's responses. For example, if only one student in a particular grade reported where he/she obtained alcohol, each category would show up as either 0% or 100%. The table indicates the sample size for each grade surveyed to help clarify the value of the data.

After that is Table 11, CSAP questions. It contains information required by communities with CSAP Grants, such as the parent attitudes regarding drinking, police response to drinking, and problems associated with drinking.

After Table 11 is Table 12, Drug Free Communities Report. It contains information required by communities with Drug Free Communities Grants, such as the perception of the risks of ATOD use, perception of parent and peer disapproval of ATOD use, past 30-day use, and average age of first use.

Next is Table 13, Youth Perceptions of Substance Use. Youth often overestimate the percentage of their peers who are using substances. Youth perceptions of the percentage of their peers who use cigarettes, alcohol, marijuana, and other illegal drugs are shown in these tables.

Finally, there are any extra questions your agency might have asked.



## What are the numbers telling you?

Review the charts and data tables presented in this report. Note your findings as you discuss the following questions.

**Which 3-5 risk factors** appear to be higher than you would want when compared to the Bach Harrison Norm?

**Which 3-5 protective factors** appear to be lower than you would want when compared to the Bach Harrison Norm?

**Which levels of 30-day drug use** are increasing and/or unacceptably high? Which substances are your students using the most? At which grades do you see unacceptable usage levels?

**Which antisocial behaviors** are increasing and/or unacceptably high? Which behaviors are your students exhibiting the most? At which grades do you see unacceptable behavior levels?

## How to identify high priority problem areas

Once you have familiarized yourself with the data, you can begin to identify priorities.

**Look across the charts** for items that stand out as either much higher or much lower than the others.

**Compare your data** with statewide, and/or national data. Differences of 5% between local and other data are probably significant.

**Prioritize problems for your area** according to the issues you've identified. Which can be realistically addressed with the funding available to your community? Which problems fit best with the prevention resources at hand?

**Determine the standards and values** held within your community. For example: Is it acceptable in your community for a percentage of high school students to drink alcohol regularly as long as that percentage is lower than the overall state rate?

## Use these data for planning.

Once priorities are established, use data to guide your prevention efforts.

**Substance use and antisocial behavior data** are excellent tools to raise awareness about the problems and promote dialogue.


**Risk and protective factor data** can be used to identify exactly where the community needs to take action.

**Promising approaches** for any prevention goal are available through resources listed on the last page of this report. These contacts are a great resource for information about programs that have been proven effective in addressing the risk factors that are high in your community, and improving the protective factors that are low.

	Sample	Priority Rate 1	Priority Rate 2	Priority Rate 3
<b>Risk Factors</b>	6th grad Fav. Attitude to Drugs (Peer/Indiv. Scale) @ 15% (8% > 8-state av.)			
<b>Protective Factors</b>	10th grad - Rewards for prosocial involvm. (School Domain) 40% (down 5% from 2 yrs ago & 16% below state av.)			
<b>30-day Substance Abuse</b>	8th grad Binge Drinking @ 13% (5% above state av.)			
<b>Antisocial Behavior</b>	12th grad - Drunk/High at School @ 21% (about same as state, but remains a priority.)			


# Substance Use and Antisocial Behavior

LIFETIME, 30 DAY & HEAVY ATOD USE  
2017 Chesterfield County, Grade 8




\* Sedatives and Other Narcotics are 2014 values.  
\*\* MTF does not collect data on those substances.

ANTISOCIAL BEHAVIOR  
2017 Chesterfield County, Grade 8




# Substance Use and Antisocial Behavior

LIFETIME, 30 DAY & HEAVY ATOD USE  
2017 Chesterfield County, Grade 10




\* Sedatives and Other Narcotics are 2014 values.  
\*\* MTF does not collect data on those substances.

ANTISOCIAL BEHAVIOR  
2017 Chesterfield County, Grade 10




# Substance Use and Antisocial Behavior

LIFETIME, 30 DAY & HEAVY ATOD USE  
2017 Chesterfield County, Grade 12



\*\* MTF does not collect data on those substances.


ANTISOCIAL BEHAVIOR  
2017 Chesterfield County, Grade 12






# Substance Use and Antisocial Behavior

LIFETIME, 30 DAY & HEAVY ATOD USE  
2017 Chesterfield County, All Grades




\* MTF only surveys grades 8, 10, and 12. \* Sedatives and Other Narcotics are 2014 values.  
\*\* MTF does not collect data on those substances.

ANTISOCIAL BEHAVIOR  
2017 Chesterfield County, All Grades




# Risk and Protective Factor Profiles

## RISK PROFILE 2017 Chesterfield County, Grade 8



\* High Risk Youth are defined as the percentage of students who have more than a specified number of risk factors operating in their lives. (6th grade: 5 or more risk factors, 7th-9th grades: 6 or more factors, 10th-12th grades: 7 or more factors)


## PROTECTIVE PROFILE 2017 Chesterfield County, Grade 8



\*\* High Protection Youth are defined as the percentage of students who have more than a specified number of protective factors operating in their lives. (6th and 7th grades: 3 or more protective factors, 8th-12th grades: 4 or more factors).


# Risk and Protective Factor Profiles

## RISK PROFILE 2017 Chesterfield County, Grade 10



\* High Risk Youth are defined as the percentage of students who have more than a specified number of risk factors operating in their lives. (6th grade: 5 or more risk factors, 7th-9th grades: 6 or more factors, 10th-12th grades: 7 or more factors)


## PROTECTIVE PROFILE 2017 Chesterfield County, Grade 10



\*\* High Protection Youth are defined as the percentage of students who have more than a specified number of protective factors operating in their lives. (6th and 7th grades: 3 or more protective factors, 8th-12th grades: 4 or more factors).


# Risk and Protective Factor Profiles

## RISK PROFILE 2017 Chesterfield County, Grade 12



\* High Risk Youth are defined as the percentage of students who have more than a specified number of risk factors operating in their lives. (6th grade: 5 or more risk factors, 7th-9th grades: 6 or more factors, 10th-12th grades: 7 or more factors)

## PROTECTIVE PROFILE 2017 Chesterfield County, Grade 12




\*\* High Protection Youth are defined as the percentage of students who have more than a specified number of protective factors operating in their lives. (6th and 7th grades: 3 or more protective factors, 8th-12th grades: 4 or more factors).




# Risk and Protective Factor Profiles

## RISK PROFILE 2017 Chesterfield County, All Grades



\* High Risk Youth are defined as the percentage of students who have more than a specified number of risk factors operating in their lives. (6th grade: 5 or more risk factors, 7th-9th grades: 6 or more factors, 10th-12th grades: 7 or more factors)

## PROTECTIVE PROFILE 2017 Chesterfield County, All Grades



\*\* High Protection Youth are defined as the percentage of students who have more than a specified number of protective factors operating in their lives. (6th and 7th grades: 3 or more protective factors, 8th-12th grades: 4 or more factors).

# Risk and Protective Scale Definitions

**Table 2. Scales that Measure the Risk and Protective Factors Shown in the Profiles**

<i>Community Domain Risk Factors</i>	
<b>Laws and Norms Favorable Toward Drug Use</b>	Research has shown that legal restrictions on alcohol and tobacco use, such as raising the legal drinking age, restricting smoking in public places, and increased taxation have been followed by decreases in consumption. Moreover, national surveys of high school seniors have shown that shifts in normative attitudes toward drug use have preceded changes in prevalence of use.
<b>Scale Questions for Laws and Norms Favorable Toward Drug Use</b>	<i>If a kid drank some beer, wine or hard liquor (for example, vodka, whiskey or gin) in your neighborhood would he or she be caught by the police?</i>
	<i>If a kid smoked marijuana in your neighborhood, would he or she be caught by the police?</i>
	<i>If a kid carried a handgun in your neighborhood, would he or she be caught by the police?</i>
	<i>How wrong would most adults (over 21) in your neighborhood think it is for kids your age to use marijuana?</i>
	<i>How wrong would most adults (over 21) in your neighborhood think it is for kids your age to drink alcohol?</i>
<b>Perceived Availability of Drugs</b>	The availability of cigarettes, alcohol, marijuana, and other illegal drugs has been related to the use of these substances by adolescents.
	<i>If you wanted to get some cigarettes, how easy would it be for you to get some?</i>
<b>Scale Questions for Perceived Availability of Drugs</b>	<i>If you wanted to get some beer, wine or hard liquor (for example, vodka, whiskey or gin), how easy would it be for you to get some?</i>
	<i>If you wanted to get some marijuana, how easy would it be for you to get some?</i>
	<i>If you wanted to get a drug like cocaine, LSD, or amphetamines, how easy would it be for you to get some?</i>
<i>Community Domain Protective Factors</i>	
<b>Rewards for Prosocial Involvement</b>	Rewards for positive participation in activities helps youth bond to the community, thus lowering their risk for substance use.
<b>Scale Questions for Rewards for Prosocial Involvement</b>	<i>My neighbors notice when I am doing a good job and let me know about it.</i>
	<i>There are people in my neighborhood who are proud of me when I do something well.</i>
	<i>There are people in my neighborhood who encourage me to do my best.</i>
<i>Family Domain Risk Factors</i>	
<b>Poor Family Management</b>	Parents' use of inconsistent and/or unusually harsh or severe punishment with their children places them at higher risk for substance use and other problem behaviors. Also, parents' failure to provide clear expectations and to monitor their children's behavior makes it more likely that they will engage in drug abuse whether or not there are family drug problems.
<b>Scale Questions for Poor Family Management</b>	<i>The rules in my family are clear.</i>
	<i>When I am not at home, one of my parents knows where I am and who I am with.</i>
	<i>If you drank some beer or wine or liquor (for example, vodka, whiskey or gin) without your parents' permission, would you be caught by your parents?</i>
	<i>My family has clear rules about alcohol use.</i>
	<i>My family has clear rules about other drug use.</i>
	<i>If you carried a handgun without your parents' permission, would you be caught by your parents?</i>
	<i>If you skipped school, would you be caught by your parents?</i>
	<i>My parents ask if I've gotten my homework done.</i>
<i>Would your parents know if you did not come home on time?</i>	

## Risk and Protective Scale Definitions

**Table 2. Scales that Measure the Risk and Protective Factors Shown in the Profiles**

<i>Family Domain Risk Factors (Continued)</i>	
<b>Family Conflict</b>	Children raised in families high in conflict, whether or not the child is directly involved in the conflict, appear at risk for both delinquency and drug use.
<b>Scale Questions for Family Conflict</b>	<i>People in my family often insult or yell at each other.</i>
	<i>We argue about the same things in my family over and over.</i>
	<i>People in my family have serious arguments.</i>
<b>Sibling Drug Use and Exposure to Adult Antisocial Behavior</b>	When children are raised in a family with a history of problem behaviors (e.g., violence or ATOD use), the children are more likely to engage in these behaviors.
<b>Scale Questions for Sibling Drug Use</b>	<i>Have any of your brothers or sisters ever:</i>
	<i>drunk beer, wine or hard liquor (for example, vodka, whiskey or gin)?</i>
	<i>smoked marijuana?</i>
	<i>smoked cigarettes?</i>
<b>Scale Questions for Exposure to Adult Antisocial Behavior</b>	<i>About how many adults (over 21) have you known personally who in the past year have:</i>
	<i>used marijuana, crack, cocaine, or other drugs?</i>
	<i>sold or dealt drugs?</i>
	<i>done other things that could get them in trouble with the police like stealing, selling stolen goods, mugging or assaulting others, etc.?</i>
	<i>gotten drunk or high?</i>
<b>Parental Attitudes Favorable Toward Antisocial Behavior and Parental Attitudes Favorable Toward Drugs</b>	In families where parents use illegal drugs, are heavy users of alcohol, or are tolerant of children's use, children are more likely to become drug abusers during adolescence. The risk is further increased if parents involve children in their own drug (or alcohol) using behavior, for example, asking the child to light the parent's cigarette or get the parent a beer from the refrigerator.
<b>Scale Questions for Parental Attitudes Favorable Toward Antisocial Behavior</b>	<i>How wrong do your parents feel it would be for YOU to steal something worth more than \$5?</i>
	<i>How wrong do your parents feel it would be for YOU to draw graffiti, write things, or draw pictures on buildings or other property (without the owner's permission)?</i>
	<i>How wrong do your parents feel it would be for YOU to pick a fight with someone?</i>
<b>Scale Questions for Parental Attitudes Favorable Toward Drug Use</b>	<i>How wrong do your parents feel it would be for YOU to drink beer, wine or hard liquor (for example, vodka, whiskey or gin) regularly?</i>
	<i>How wrong do your parents feel it would be for YOU to smoke cigarettes?</i>
	<i>How wrong do your parents feel it would be for YOU to smoke marijuana?</i>
<i>Family Domain Protective Factors</i>	
<b>Family Attachment</b>	Young people who feel that they are a valued part of their family are less likely to engage in substance use and other problem behaviors.
<b>Scale Questions for Family Attachment</b>	<i>Do you feel very close to your mother?</i>
	<i>Do you share your thoughts and feelings with your mother?</i>
	<i>Do you share your thoughts and feelings with your father?</i>
	<i>Do you feel very close to your father?</i>

## Risk and Protective Scale Definitions

**Table 2. Scales that Measure the Risk and Protective Factors Shown in the Profiles**

<i>Family Domain Protective Factors (Continued)</i>	
<b>Opportunities for Prosocial Involvement</b>	Young people who are exposed to more opportunities to participate meaningfully in the responsibilities and activities of the family are less likely to engage in drug use and other problem behaviors.
<b>Scale Questions for Opportunities for Prosocial Involvement</b>	<i>If I had a personal problem, I could ask my mom or dad for help.</i>
	<i>My parents give me lots of chances to do fun things with them.</i>
	<i>My parents ask me what I think before most family decisions affecting me are made</i>
<b>Rewards for Prosocial Involvement</b>	When parents, siblings, and other family members praise, encourage, and attend to things done well by their child, children are less likely to engage in substance use and problem behaviors.
<b>Scale Questions for Rewards for Prosocial Involvement</b>	<i>My parents notice when I am doing a good job and let me know about it.</i>
	<i>How often do your parents tell you they're proud of you for something you've done?</i>
	<i>Do you enjoy spending time with your mother?</i>
	<i>Do you enjoy spending time with your father?</i>
<i>School Domain Risk Factors</i>	
<b>Academic Failure</b>	Beginning in the late elementary grades (grades 4-6) academic failure increases the risk of both drug abuse and delinquency. It appears that the experience of failure itself, for whatever reasons, increases the risk of problem behaviors.
<b>Scale Questions for Academic Failure</b>	<i>Putting them all together, what were your grades like last year?</i>
	<i>Are your school grades better than the grades of most students in your class?</i>
<b>Low Commitment to School</b>	Surveys of high school seniors have shown that the use of drugs is significantly lower among students who expect to attend college than among those who do not. Factors such as liking school, spending time on homework, and perceiving the coursework as relevant are also negatively related to drug use.
<b>Scale Questions for Low Commitment to School</b>	<i>During the LAST FOUR WEEKS how many whole days of school have you missed because you skipped or "cut"?</i>
	<i>How often do you feel that the school work you are assigned is meaningful and important?</i>
	<i>How interesting are most of your courses to you?</i>
	<i>How important do you think the things you are learning in school are going to be for your later life?</i>
	<i>Now, thinking back over the past year in school, how often did you enjoy being in school?</i>
	<i>Now, thinking back over the past year in school, how often did you hate being in school?</i>
<i>Now, thinking back over the past year in school, how often did you try to do your best work in school?</i>	
<i>School Domain Protective Factors</i>	
<b>Opportunities for Prosocial Involvement</b>	When young people are given more opportunities to participate meaningfully in important activities at school, they are less likely to engage in drug use and other problem behaviors.
<b>Scale Questions for Opportunities for Prosocial Involvement</b>	<i>In my school, students have lots of chances to help decide things like class activities and rules.</i>
	<i>There are lots of chances for students in my school to get involved in sports, clubs, or other school activities outside of class.</i>
	<i>There are lots of chances for students in my school to talk with a teacher one-on-one.</i>
	<i>I have lots of chances to be part of class discussions or activities.</i>



## Risk and Protective Scale Definitions

**Table 2. Scales that Measure the Risk and Protective Factors Shown in the Profiles**

<i>School Domain Protective Factors (Continued)</i>	
<b>Rewards for Prosocial Involvement</b>	When young people are recognized and rewarded for their contributions at school, they are less likely to be involved in substance use and other problem behaviors.
<b>Scale Questions for Rewards for Prosocial Involvement</b>	<i>My teachers notice when I am doing a good job and lets me know about it.</i>
	<i>The school lets my parents know when I have done something well.</i>
	<i>My teachers praise me when I work hard in school.</i>
<i>Peer-Individual Risk Factors</i>	
<b>Early Initiation of Antisocial Behavior and Early Initiation of Drug Use</b>	Early onset of drug use predicts misuse of drugs. The earlier the onset of any drug use, the greater the involvement in other drug use and the greater frequency of use. Onset of drug use prior to the age of 15 is a consistent predictor of drug abuse, and a later age of onset of drug use has been shown to predict lower drug involvement and a greater probability of discontinuation of use.
<b>Scale Questions for Early Initiation of Antisocial Behavior</b>	<i>How old were you when you first...</i>
	<i>got suspended from school?</i>
	<i>got arrested?</i>
	<i>carried a handgun?</i>
	<i>attacked someone with the idea of seriously hurting them?</i>
<b>Scale Questions for Early Initiation of Drug Use</b>	<i>How old were you when you first...</i>
	<i>smoked a cigarette, even just a puff?</i>
	<i>had more than a sip or two of beer, nine or hard liquor (for example, vodka, whiskey or gin)?</i>
	<i>began drinking alcoholic beverages regularly, that is, at least once or twice a month?</i>
	<i>smoked marijuana?</i>
<b>Attitudes Favorable Toward Antisocial Behavior and Attitudes Favorable Toward Drug Use</b>	During the elementary school years, most children express anti-drug, anti-crime, and pro-social attitudes and have difficulty imagining why people use drugs or engage in antisocial behaviors. However, in middle school, as more youth are exposed to others who use drugs and engage in antisocial behavior, their attitudes often shift toward greater acceptance of these behaviors. Youth who express positive attitudes toward drug use and antisocial behavior are more likely to engage in a variety of problem behaviors, including drug use.
<b>Scale Questions for Attitudes Favorable Toward Antisocial Behavior</b>	<i>How wrong do you think it is for someone your age to...</i>
	<i>take a handgun to school?</i>
	<i>steal anything worth more than \$5?</i>
	<i>pick a fight with someone?</i>
	<i>attack someone with the idea of seriously hurting them?</i>
<b>Scale Questions for Attitudes Favorable Toward Drug Use</b>	<i>stay away from school all day when their parents think they are at school?</i>
	<i>How wrong do you think it is for someone your age to...</i>
	<i>drink beer, nine or hard liquor (for example, vodka, whiskey or gin) regularly?</i>
	<i>smoke cigarettes?</i>
	<i>smoke marijuana?</i>
	<i>use LSD, cocaine, amphetamines or another illegal drug?</i>

## Risk and Protective Scale Definitions

**Table 2. Scales that Measure the Risk and Protective Factors Shown in the Profiles**

<i>Peer-Individual Risk Factors (Continued)</i>	
<b>Perceived Risk of Drug Use</b>	Young people who do not perceive drug use to be risky are far more likely to engage in drug use.
<b>Scale Questions for Perceived Risk of Drug Use</b>	<i>How much do you think people risk harming themselves (physically or in other ways) if they:</i>
	<i>smoke one or more packs of cigarettes per day?</i>
	<i>try marijuana once or twice?</i>
	<i>smoke marijuana regularly?</i>
	<i>take one or two drinks of an alcoholic beverage (beer, wine, or liquor) nearly every day?</i>
<b>Interaction with Antisocial Peers</b>	Young people who associate with peers who engage in problem behaviors are at higher risk for engaging in antisocial behavior themselves.
<b>Scale Questions for Interaction with Antisocial Peers</b>	<i>Think of your four best friends (the friends you feel closest to). In the past year (12 months), how many of your best friends have:</i>
	<i>been suspended from school?</i>
	<i>carried a handgun?</i>
	<i>sold illegal drugs?</i>
	<i>stolen or tried to steal a motor vehicle such as a car or motorcycle?</i>
	<i>been arrested?</i>
	<i>dropped out of school?</i>
<b>Friends' Use of Drugs</b>	Young people who associate with peers who engage in alcohol or substance abuse are much more likely to engage in the same behavior. Peer drug use has consistently been found to be among the strongest predictors of substance use among youth. Even when young people come from well-managed families and do not experience other risk factors, spending time with friends who use drugs greatly increases the risk of that problem developing.
<b>Scale Questions for Friends' Use of Drugs</b>	<i>Think of your four best friends (the friends you feel closest to). In the past year (12 months), how many of your best friends have:</i>
	<i>smoked cigarettes?</i>
	<i>tried beer, wine or hard liquor (for example, vodka, whiskey or gin) when their parents didn't know about it?</i>
	<i>used marijuana?</i>
	<i>used LSD, cocaine, amphetamines, or other illegal drugs?</i>
<b>Rewards for Antisocial Behavior</b>	Young people who receive rewards for their antisocial behavior are at higher risk for engaging further in antisocial behavior and substance use.
<b>Scale Questions for Rewards for Antisocial Behavior</b>	<i>What are the chances you would be seen as cool if you:</i>
	<i>smoked cigarettes?</i>
	<i>began drinking alcoholic beverages regularly, that is, at least once or twice a month?</i>
	<i>smoked marijuana?</i>
<b>Depressive Symptoms</b>	Young people who are depressed are overrepresented in the criminal justice system and are more likely to use drugs. Survey research and other studies have shown a link between depression and youth problem behaviors.
<b>Scale Questions for Depressive Symptoms</b>	<i>Sometimes I think that life is not worth it.</i>
	<i>At times I think I am no good at all.</i>
	<i>All in all, I am inclined to think that I am a failure.</i>
	<i>In the past year, have you felt depressed or sad MOST days, even if you felt OK sometimes?</i>

## Risk and Protective Scale Definitions

**Table 2. Scales that Measure the Risk and Protective Factors Shown in the Profiles**

<i>Peer-Individual Risk Factors (Continued)</i>	
<b>Gang Involvement</b>	Youth who belong to gangs are more at risk for antisocial behavior and drug use.
<b>Scale Question for Gang Involvement</b>	<i>Have you ever belonged to a gang?</i>
<i>Peer-Individual Protective Factors</i>	
<b>Religiosity</b>	Young people who regularly attend religious services are less likely to engage in problem behaviors.
<b>Scale Question for Religiosity</b>	<i>How often do you attend religious services or activities?</i>
<b>Belief in the Moral Order</b>	Young people who have a belief in what is “right” or “wrong” are less likely to use drugs.
<b>Scale Questions for Belief in the Moral Order</b>	<i>It is all right to beat up people if they start the fight.</i>
	<i>It is important to be honest with your parents, even if they become upset or you get punished.</i>
	<i>I think it is okay to take something without asking if you can get away with it.</i>
	<i>I think sometimes it’s okay to cheat at school.</i>
<b>Interaction with Prosocial Peers</b>	Young people who associate with peers who engage in prosocial behavior are more protected from engaging in antisocial behavior and substance use.
<b>Scale Questions for Interaction with Antisocial Peers</b>	<i>Think of your four best friends (the friends you feel closest to). In the past year (12 months), how many of your best friends have:</i>
	<i>participated in clubs, organizations or activities at school?</i>
	<i>made a commitment to stay drug-free?</i>
	<i>tried to do well in school?</i>
	<i>liked school?</i>
	<i>regularly attended religious services?</i>
<b>Prosocial Involvement</b>	Participation in positive school and community activities helps provide protection for youth.
<b>Scale Questions for Prosocial Involvement</b>	<i>How many times in the past year (12 months) have you:</i>
	<i>participated in clubs, organizations or activities at school?</i>
	<i>done extra work on your own for school?</i>
	<i>volunteered to do community service?</i>
<b>Rewards for Prosocial Involvement</b>	Young people who are rewarded for working hard in school and the community are less likely to engage in problem behavior.
<b>Scale Questions for Rewards for Prosocial Involvement</b>	<i>What are the chances you would be seen as cool if you:</i>
	<i>worked hard at school?</i>
	<i>defended someone who was being verbally abused at school?</i>
	<i>regularly volunteered to do community service?</i>

# Data Tables

**Table 3. Number of Students Who Completed the Survey**

Number of Youth	Grade 8				Grade 10				Grade 12				Total			
	2014	2016	2017	MTF †	2014	2016	2017	MTF †	2014	2016	2017	MTF †	2014	2016	2017	MTF †
	1373	1096	1940	†	1161	1266	1254	†	1211	1152	947	†	3745	3514	4141	†

**Table 4. Percentage of Students Who Used ATODs During Their Lifetime**

Substance	Grade 8				Grade 10				Grade 12				Total			
	2014	2016	2017	MTF †	2014	2016	2017	MTF †	2014	2016	2017	MTF †	2014	2016	2017	MTF †
Alcohol	27.4	21.7	13.2	22.8	49.1	36.8	29.0	43.4	68.5	54.3	51.2	61.2	47.5	37.6	26.8	41.9
Cigarettes	12.5	9.0	2.9	9.8	21.5	14.5	8.0	17.5	36.3	27.3	22.7	28.3	23.0	16.9	8.9	18.2
E-Cigarettes	n/a	n/a	4.6	17.5	n/a	n/a	15.0	29.0	n/a	n/a	29.0	33.8	n/a	n/a	13.3	26.6
Chewing Tobacco	3.7	4.1	1.3	6.9	10.4	7.5	3.2	10.2	16.5	13.4	9.8	14.2	10.0	8.3	3.8	10.3
Marijuana	10.6	10.5	2.8	12.8	27.1	21.2	13.5	29.7	46.4	37.0	35.4	44.5	27.3	22.8	13.5	28.6
Inhalants	8.9	6.6	4.6	7.7	6.1	3.6	4.1	6.6	6.1	5.0	2.4	5.0	7.1	5.0	4.0	6.5
Hallucinogens	1.4	1.6	0.5	1.9	3.6	3.1	1.4	4.4	7.8	8.3	5.0	6.7	4.1	4.3	1.8	4.3
Cocaine	1.0	0.7	0.3	1.4	1.6	0.7	0.8	2.1	4.0	4.2	3.2	3.7	2.2	1.8	1.1	2.3
Methamphetamines	0.2	0.6	0.1	0.6	0.7	0.5	0.2	0.7	1.5	1.3	0.8	1.2	0.8	0.8	0.3	0.8
Amphetamines	3.6	3.9	1.1	5.7	9.5	5.2	2.4	8.8	15.7	10.6	8.2	10.0	9.4	6.5	3.1	8.1
Sedatives *	6.1	7.6	2.0	5.5	7.3	6.3	2.4	8.1	10.0	6.2	3.8	5.2	7.7	6.7	2.5	6.7
Tranquilizers	1.2	2.7	0.7	3.0	4.4	4.2	2.8	6.1	8.2	9.1	6.8	7.6	4.5	5.3	2.7	5.5
Heroin	0.5	0.4	0.1	0.5	0.7	0.7	0.3	0.6	0.6	0.8	0.5	0.7	0.6	0.6	0.2	0.6
Other Narcotics *	1.9	2.2	0.8	2.3	6.9	3.4	1.9	6.8	12.4	9.5	6.2	n/a	6.9	5.0	2.3	6.1
Ecstasy	1.0	0.6	0.5	1.7	2.2	1.5	0.7	2.8	5.3	3.4	2.0	4.9	2.8	1.8	0.9	3.1
Synthetic Marijuana **	n/a	1.6	0.6	n/a	n/a	2.2	0.4	n/a	n/a	3.6	2.6	n/a	n/a	2.5	1.0	n/a

† See the Monitoring The Future (MTF) website ( [www.monitoringthefuture.org](http://www.monitoringthefuture.org) ). MTF only surveys grades 8, 10 and 12.

\* The Substances in Blue are 2014 values.

\*\* MTF does not collect data on that substance.

# Data Tables

**Table 5. Percentage of Students Who Used ATODs During The Past 30 Days**

Substance	Grade 8				Grade 10				Grade 12				Total			
	2014	2016	2017	MTF †	2014	2016	2017	MTF †	2014	2016	2017	MTF †	2014	2016	2017	MTF †
Alcohol	10.0	8.7	4.9	7.3	26.1	18.3	14.4	19.9	41.5	36.2	32.6	33.2	25.2	21.0	14.2	19.8
Cigarettes	2.7	1.6	0.9	2.6	7.4	4.2	2.7	4.9	12.9	9.0	9.6	10.5	7.5	4.9	3.4	5.9
E-Cigarettes	n/a	n/a	2.7	6.2	n/a	n/a	8.2	11.0	n/a	n/a	17.9	12.5	n/a	n/a	7.8	9.9
Chewing Tobacco	1.5	1.4	1.2	2.5	5.6	3.3	2.3	3.5	7.4	6.7	5.6	6.6	4.7	3.8	2.5	4.1
Marijuana	4.9	4.9	1.3	5.4	14.6	9.9	6.0	14.0	24.7	21.8	19.5	22.5	14.4	12.1	6.9	13.7
Inhalants	2.5	2.4	1.9	1.8	1.4	0.8	0.9	1.0	1.0	1.3	0.3	0.8	1.7	1.5	1.2	1.2
Hallucinogens	0.2	0.2	0.2	0.6	1.4	1.2	0.3	0.9	2.9	3.6	1.1	1.4	1.5	1.6	0.4	1.0
Cocaine	0.5	0.3	0.1	0.3	0.3	0.4	0.3	0.4	1.4	1.1	0.6	0.9	0.7	0.6	0.3	0.5
Methamphetamines	0.2	0.2	0.0	0.3	0.3	0.1	0.0	0.2	0.5	0.4	0.0	0.3	0.3	0.2	0.0	2.5
Amphetamines	1.8	1.6	0.2	1.7	4.1	2.0	0.9	2.7	5.5	3.8	3.2	3.0	3.7	2.4	1.1	0.3
Sedatives *	2.6	2.6	0.6	1.6	3.0	2.8	1.1	2.3	3.9	2.3	0.8	1.5	3.2	2.6	0.8	2.0
Tranquilizers	0.5	1.0	0.4	0.8	1.5	1.6	1.1	1.5		3.2	1.2	1.9	1.6	1.9	0.8	1.4
Heroin	0.2	0.3	0.0	0.2	0.3	0.5	0.0	0.2	0.1	0.1	0.0	0.2	0.2	0.3	0.0	0.2
Other Narcotics *	0.7	1.2	0.2	0.7	2.9	1.6	0.4	1.7	3.4	3.0	1.2	1.7	2.3	1.9	0.5	1.4
Ecstasy	0.4	0.2	0.2	0.3	0.4	0.5	0.0	0.5	0.9	0.9	0.2	0.9	0.6	0.5	0.1	0.6
Synthetic Marijuana **	n/a	0.6	0.2	n/a	n/a	1.0	0.0	0.0	n/a	0.9	0.6	0.0	n/a	0.8	0.2	0.0
Energy Drinks **	n/a	22.3	11.7	n/a	n/a	20.9	14.8	n/a	n/a	24.8	21.6	n/a	n/a	22.6	14.9	n/a
Caffeine Pills **	n/a	1.8	0.7	n/a	n/a	2.4	0.4	n/a	n/a	2.3	2.0	n/a	n/a	2.2	0.9	n/a

**Table 6. Percentage of Students With Problem ATOD Use**

Problem Use	2014	2016	2017	MTF †	2014	2016	2017	MTF †	2014	2016	2017	MTF †	2014	2016	2017	MTF †
Binge Drinking (past 2 weeks)	3.3	4.3	2.3	3.4	12.5	10.1	7.4	9.7	20.9	22.7	20.3	15.5	11.9	12.3	8.0	9.4
1/2 Pack of Cigarettes per Day	0.3	0.1	0.4	0.3	1.3	0.6	0.8	0.6	3.1	1.5	1.8	1.8	1.5	0.7	0.8	0.9
<b>Alcohol and Driving in the Past 30 Days</b>	2014	2016	2017	BH Norm	2014	2016	2017	BH Norm	2014	2016	2017	BH Norm	2014	2016	2017	BH Norm
Drinking and Driving	0.6	1.6	1.6	5.6	2.5	2.5	1.3	5.3	7.2	6.3	4.3	11.8	3.3	3.4	2.1	7.5
Riding with a Drinking Driver	17.4	17.5	12.1	22.3	16.3	16.6	13.6	24.0	16.5	13.5	12.2	24.1	16.8	15.9	12.6	23.5

† See the Monitoring The Future (MTF) website ( [www.monitoringthefuture.org](http://www.monitoringthefuture.org) ). MTF only surveys grades 8, 10 and 12.

\* The Substances in Blue are 2014 values.

\*\* MTF does not collect data on that substance.

# Data Tables, Supplement 1

## Supplement 1. ATOD Questions

Cigarettes	"Have you ever smoked cigarettes?" And "How frequently have you smoked cigarettes during the past 30 days?"
Chewing Tobacco	"Have you ever used smokeless tobacco (chew, snuff, plug, dipping tobacco, or chewing tobacco)? " and "How frequently have you used smokeless tobacco during the past 30 days?"
<b>On how many occasions (if any) have you:</b>	
Alcohol	had alcoholic beverages (beer, wine or hard liquor) to drink in your lifetime (or during the past 30 days) -- more than just a few sips?
E-Cigarettes	used electronic cigarettes, e-cigarettes, vape pens, or e-hookahs in your lifetime (or during the past 30 days)?
Marijuana	used marijuana (grass, pot) or hashish (hash, hash oil) in your lifetime (or during the past 30 days)?
Inhalants	sniffed glue, breathed the contents of an aerosol spray can, or inhaled other gases or sprays, in order to get high in your lifetime (or during the past 30 days)?
Hallucinogens	used LSD (acid) or other hallucinogens (like PCP, mescaline, peyote, "shrooms" or psilocybin) in your lifetime (or during the past 30 days)?
Cocaine	used cocaine (like cocaine powder) or "crack" (cocaine in chunk or rock form) in your lifetime (or during the past 30 days)?
Methamphetamines	used methamphetamines (meth, speed, crank, crystal meth) in your lifetime (or during the past 30 days)?
Amphetamines	used prescription stimulants or amphetamines (such as Adderall, Ritalin, or Dexedrine) without a doctor telling you to take them, in your lifetime (or during the past 30 days)?
Sedatives	used prescription sedatives including barbiturates or sleeping pills (such as phenobarbital, Tuinal, Seconal, Ambien, Lunesta, or Sonata) without a doctor telling you to take them, in your lifetime (or during the past 30 days)?
Tranquilizers	used prescription tranquilizers (such as Librium, Valium, Xanax, Ativan, Soma, or Klonopin) without a doctor telling you to take them, in your lifetime (or during the past 30 days)?
Heroin	used heroin in your lifetime (or during the past 30 days)?
Other Narcotics	used narcotic prescription drugs (such as OxyContin, methadone, morphine, codeine, Demerol, Vicodin, or Percocet) without a doctor telling you to take them, in your lifetime (or during the past 30 days)?
Ecstasy	used MDMA (X,E, or ecstasy) in your lifetime (or during the past 30 days)?
Synthetic Marijuana	used "synthetic marijuana" ("K2", "Spice") to get high in your lifetime (or during the past 30 days)?
Energy Drinks	drank energy drinks with caffeine (like Red Bull, Monster, Rockstar, or 5-Hour-Energy) during the past 30 days?
Caffeine Pills	used caffeine pills (No-Doz, Vivarin, Dexatrim) during the past 30 days?
<b>Problem Use</b>	
Binge Drinking (past 2 weeks)	Think back over the last two weeks. How many times have you had five or more alcoholic drinks in a row? ( A "drink" is a glass of wine, a bottle of beer, a wine cooler, a shot glass of liquor, or a mixed drink.)
1/2 Pack of Cigarettes per Day	How frequently have you smoked cigarettes during the past 30 days?
<b>Alcohol and Driving in the Past 30 Days</b>	
Drinking and Driving	During the past 30 days, how many times did you DRIVE a car or other vehicle when you had been drinking alcohol?
Riding with a Drinking Driver	During the past 30 days, how many times did you RIDE in a car or other vehicle driven by someone who had been drinking alcohol?





# Data Tables

**Table 7. Percentage of Students With Antisocial Behavior in the Past Year**

How many times in the past year (12 months) have you: (One or more times)	Grade 8				Grade 10				Grade 12				Total			
	2014	2016	2017	BH Norm	2014	2016	2017	BH Norm	2014	2016	2017	BH Norm	2014	2016	2017	BH Norm
Been Suspended from School	10.3	10.5	7.8	13.4	8.7	6.5	7.0	11.2	8.2	6.7	7.6	8.5	9.1	7.9	7.5	11.1
Been Drunk or High at School	5.0	5.0	2.1	7.8	12.6	7.9	5.8	14.7	19.5	14.2	14.0	17.3	12.0	9.0	6.0	13.2
Sold Illegal Drugs	2.2	2.0	0.8	3.1	5.3	4.9	2.5	7.2	8.2	8.1	5.4	8.6	5.1	5.0	2.4	6.3
Stolen or Tried to Steal a Motor Vehicle	0.7	1.1	0.7	2.2	1.7	1.4	2.2	2.7	1.8	1.7	1.7	2.0	1.4	1.4	1.4	2.3
Been Arrested	2.2	1.4	0.8	4.8	3.2	1.7	2.1	6.0	4.3	3.3	4.2	5.8	3.2	2.1	2.0	5.5
Attacked Someone with the Idea of Seriously Hurting Them	8.8	8.1	6.4	12.9	7.5	5.8	3.6	11.8	8.3	5.3	6.0	9.6	8.3	6.4	5.4	11.5
Carried a Handgun	5.0	4.0	5.0	5.4	5.2	3.6	5.3	5.5	6.5	5.4	5.1	5.5	5.5	4.3	5.1	5.5
Carried a Handgun to School	0.3	0.8	0.6	0.9	0.7	1.0	1.3	1.2	0.8	1.1	1.8	1.2	0.6	0.9	1.1	1.1

# Data Tables

**Table 8. Percentage of Students Reporting Protection**

Protective Factors	Grade 8				Grade 10				Grade 12				Total			
	2014	2016	2017	BH Norm	2014	2016	2017	BH Norm	2014	2016	2017	BH Norm	2014	2016	2017	BH Norm
<b>Community Domain</b>																
Rewards for Prosocial Involvement	37.3	33.4	34.4	52.1	33.9	27.8	29.1	45.2	33.0	28.3	19.2	44.5	34.8	29.8	29.4	47.4
<b>Family Domain</b>																
Family Attachment	52.9	55.1	64.7	54.8	55.8	58.4	63.5	56.8	53.5	60.8	51.3	57.7	54.0	58.1	61.3	56.4
Opportunity for Prosocial Involvement	65.4	60.1	65.0	62.5	56.2	55.1	59.8	56.2	55.3	57.0	51.3	56.2	59.2	57.4	60.3	58.3
Rewards for Prosocial Involvement	43.9	46.0	53.1	48.8	46.1	50.5	58.2	54.3	43.8	51.1	45.8	54.0	44.6	49.2	53.0	52.4
<b>School Domain</b>																
Opportunity for Prosocial Involvement	56.3	52.0	57.8	62.1	48.6	52.9	57.5	63.3	51.1	49.4	51.7	64.9	52.3	51.5	56.3	63.4
Rewards for Prosocial Involvement	43.3	41.6	45.6	56.2	54.1	58.3	57.1	68.0	52.1	55.5	48.9	69.8	49.5	52.1	49.9	64.6
<b>Peer-Individual Domain</b>																
Religiosity	56.6	51.8	49.6	53.7	51.3	47.5	46.2	48.4	46.1	41.0	37.3	42.9	51.6	46.8	45.8	48.4
Belief in the Moral Order	72.8	71.7	83.4	65.8	60.6	67.0	75.9	54.6	57.1	64.9	67.5	55.6	63.9	67.9	77.4	58.7
Interaction with Prosocial Peers *	44.9	42.9	43.8	54.9	45.7	41.0	46.8	55.4	45.6	39.9	42.8	57.3	45.4	41.3	44.5	55.8
Prosocial Involvement	64.6	62.7	62.7	58.1	59.9	56.8	58.1	58.2	58.4	50.3	52.2	58.9	61.2	56.6	58.9	58.4
Rewards for Prosocial Involvement	44.2	48.8	49.2	50.9	54.2	57.9	59.9	59.9	50.6	51.4	51.8	52.2	49.4	52.9	53.2	54.4
<b>High Protection</b>																
High Protection Youth **	62.3	60.4	49.2	43.3	60.7	54.2	51.1	47.3	59.3	54.5	41.7	47.3	60.9	56.2	48.1	45.9

\*\* High Protection Youth are defined as the percentage of students who have more than a specified number of protective factors operating in their lives. (6th and 7th grades: 3 or more protective factors, 8th-12th grades: 4 or more factors).

# Data Tables

**Table 9. Percentage of Students Reporting Risk**

Risk Factors	Grade 8				Grade 10				Grade 12				Total			
	2014	2016	2017	BH Norm	2014	2016	2017	BH Norm	2014	2016	2017	BH Norm	2014	2016	2017	BH Norm
<b>Community Domain</b>																
Laws & Norms Favor Drug Use	37.8	41.9	32.7	40.0	40.2	48.5	39.8	42.3	54.8	58.2	63.3	48.1	44.2	49.5	41.8	43.4
Perceived Availability of Drugs	31.8	30.3	14.3	36.9	32.9	29.0	16.0	38.6	37.4	36.2	27.6	41.0	34.0	31.8	17.8	38.8
<b>Family Domain</b>																
Poor Family Management	37.8	47.8	32.3	40.4	43.2	42.7	28.4	40.0	46.8	46.6	37.8	41.2	42.4	45.6	32.3	40.5
Family Conflict	35.7	35.7	33.6	35.3	42.2	38.0	33.0	39.9	40.7	36.2	42.3	38.0	39.4	36.7	35.4	37.8
Sibling Drug Use	38.1	41.7	25.4	43.3	33.8	29.9	21.0	39.1	48.7	41.4	34.6	47.1	40.2	37.5	26.1	43.1
Exposure to Adult Antisocial Behavior	38.8	39.1	20.1	39.9	45.4	40.3	21.2	46.6	47.5	37.5	29.0	42.9	43.7	39.0	22.5	43.2
Parent Attitudes Favor Antisocial Behavior	47.2	46.3	30.6	49.1	49.5	42.9	37.2	53.5	52.2	43.3	37.1	52.9	49.6	44.2	34.1	51.8
Parent Attitudes Favor Drug Use	24.9	24.3	14.1	23.7	41.0	36.8	27.1	39.6	45.9	41.5	34.8	40.3	36.8	34.3	22.9	34.3
<b>School Domain</b>																
Academic Failure	39.9	39.7	30.2	37.2	40.9	39.5	34.6	39.8	44.8	44.8	48.8	37.9	41.8	41.3	35.8	38.3
Low Commitment to School	49.9	56.3	52.4	45.1	50.3	53.4	45.0	41.1	53.9	60.5	52.2	42.1	51.3	56.6	50.0	42.8
<b>Peer-Individual Domain</b>																
Early Initiation of Antisocial Behavior	27.2	28.6	15.0	32.2	31.2	24.8	17.1	34.2	36.2	29.7	23.7	34.2	31.3	27.6	17.7	33.6
Early Initiation of Drug Use	24.5	23.4	9.2	31.9	26.0	20.5	11.5	32.8	37.4	30.4	26.9	38.0	29.1	24.6	14.0	34.2
Attitudes Favorable to Antisocial Behavior	32.4	29.9	18.4	34.7	36.9	31.6	28.1	40.8	40.6	33.1	28.1	39.0	36.5	31.5	23.7	38.1
Attitudes Favorable to Drug Use	31.7	27.4	14.9	33.0	46.5	38.0	28.8	45.2	50.8	44.7	37.8	46.9	42.4	36.7	24.6	41.6
Perceived Risk of Drug Use	46.0	51.7	46.0	37.9	63.0	64.3	61.8	51.9	64.2	64.7	63.7	47.4	57.2	60.3	55.0	45.6
Interaction with Antisocial Peers	24.8	26.7	17.9	30.0	30.2	23.4	20.3	31.3	32.0	28.2	27.3	29.6	28.8	26.0	20.8	30.3
Friend's Use of Drugs	29.6	28.5	16.0	39.2	35.7	27.8	21.2	40.4	37.8	32.7	28.3	38.5	34.2	29.6	20.5	39.4
Rewards for Antisocial Behavior	41.1	32.9	19.5	35.1	51.7	42.5	34.3	46.7	56.8	47.2	42.1	51.5	49.5	41.0	29.4	44.1
Depressive Symptoms	39.6	40.0	31.9	34.8	44.2	40.8	31.9	37.8	40.4	39.7	45.7	33.4	41.3	40.2	35.0	35.4
Gang Involvement	3.8	4.7	2.8	6.9	4.1	4.0	2.9	5.9	4.4	4.2	2.1	5.2	4.1	4.3	2.7	6.0
<b>High Risk</b>																
High Risk Youth *	43.0	44.0	17.0	36.7	47.5	36.3	18.7	40.0	55.8	44.7	30.7	40.8	48.5	41.4	20.6	39.1

\* High Risk Youth are defined as the percentage of students who have more than a specified number of risk factors operating in their lives. (6th grade: 5 or more risk factors, 7th-9th grades: 6 or more factors, 10th-12th grades: 7 or more factors).

# Student Alcohol Table

**Table 10. Sources and Places of Student Alcohol Use**

If you drank alcohol (not just a sip or taste) in the past year (12 months), how did you get it?	Grade 8			Grade 10			Grade 12			Total		
	2014	2016	2017	2014	2016	2017	2014	2016	2017	2014	2016	2017
<b>Sample size *</b>	<b>288</b>	<b>183</b>	<b>144</b>	<b>517</b>	<b>389</b>	<b>240</b>	<b>747</b>	<b>527</b>	<b>306</b>	<b>1,552</b>	<b>1,099</b>	<b>690</b>
I bought it myself from a store.	0.7	4.9	5.6	3.9	5.9	7.1	12.2	13.9	15.7	7.3	9.6	10.6
I got it at a party.	35.8	36.6	40.3	58.4	52.7	52.5	74.2	69.6	66.0	61.8	58.1	55.9
I gave someone else money to buy it for me.	8.7	15.8	9.0	24.0	20.8	22.5	46.3	47.6	38.9	31.9	32.8	27.0
I got it from someone I know age 21 or older.	43.8	43.2	35.4	52.2	46.5	48.3	68.5	66.6	54.9	58.5	55.6	48.6
I got it from someone I know under age 21.	21.5	26.8	21.5	42.4	37.8	36.7	43.6	39.7	41.8	39.1	36.9	35.8
I got it from a family member or relative other than my parents.	43.4	43.2	36.8	42.9	38.6	45.0	41.0	38.1	39.5	42.1	39.1	40.9
I got it from home with my parents' permission.	42.0	49.7	53.5	42.6	42.7	46.7	39.6	41.7	46.1	41.0	43.4	47.8
I got it from home without my parents' permission.	44.8	39.3	21.5	47.4	39.8	33.8	40.0	31.1	34.6	43.4	35.6	31.6
I got it at work.	14.9	3.8	4.2	12.0	3.1	5.4	13.5	8.2	9.8	13.3	5.6	7.1
I bought it over the internet.	474.3	2.2	5.6	222.8	2.8	5.0	160.9	2.8	3.6	239.7	2.7	4.5
I got it in another way.	474.3	13.1	12.5	222.8	10.8	11.7	160.9	6.8	11.1	239.7	9.3	11.6

During the past year (12 months) did you drink alcohol at any of the following places?	Grade 8			Grade 10			Grade 12			Total		
	2014	2016	2017	2014	2016	2017	2014	2016	2017	2014	2016	2017
<b>Sample size *</b>	<b>313</b>	<b>210</b>	<b>175</b>	<b>523</b>	<b>384</b>	<b>238</b>	<b>736</b>	<b>503</b>	<b>299</b>	<b>1,572</b>	<b>1,097</b>	<b>712</b>
At my home or someone else's home without any parent permission.	49.2	43.8	30.9	66.2	57.3	55.5	67.8	62.0	59.2	63.5	56.9	51.0
At my home with my parent's permission.	54.3	59.5	66.9	50.1	50.3	55.5	49.7	51.1	56.5	50.8	52.4	58.7
At someone else's home with their parent's permission.	16.9	18.1	14.3	26.6	26.6	25.2	44.0	52.1	47.2	32.8	36.6	31.7
At an open area like a park, beach, or back road.	13.1	16.2	14.9	25.6	21.1	20.6	28.0	32.0	28.8	24.2	25.2	22.6
At public events such as a sporting event, festival, or concert.	7.7	10.5	12.0	16.4	18.5	17.6	26.0	35.4	30.4	19.1	24.7	21.6
At a restaurant, bar, or a nightclub.	9.3	6.7	8.0	11.5	9.6	11.3	16.6	16.9	18.7	13.4	12.4	13.6
In a car.	10.5	12.9	9.7	20.1	13.5	16.4	29.2	31.0	23.4	22.5	21.4	17.7
At a school dance, a game, or other event.	6.4	5.2	8.6	10.7	9.1	13.0	17.3	18.1	23.4	12.9	12.5	16.3
At school during the day.	4.8	6.7	7.4	7.8	5.7	10.1	6.5	6.8	6.7	6.6	6.4	8.0
Near school.	5.1	5.2	7.4	8.0	5.2	10.1	11.7	9.7	12.4	9.2	7.3	10.4
In another place.	19.8	15.2	22.9	20.5	15.1	19.3	23.8	18.9	22.1	21.9	16.9	21.3

\* Sample size represents the number of youth who answered the question, not including students reporting no use in the past year. In the case of smaller sample sizes, caution should be exercised before generalizing results and yearly trends to the entire community.

# Data Tables

**Table 11. CSAP Questions**

Question	Response	Grade 8		Grade 10		Grade 12		Total	
		Number	Percent	Number	Percent	Number	Percent	Number	Percent
<i>For the following for questions, during the past 12 months, have you talked with at least one of your parents (by parents, we mean either your biological parents, adoptive parents, stepparents, foster parents, or other adult caregivers whether or not they live with you) about :</i>									
The dangers of underage drinking?	Yes	796	62.4	530	61.6	340	54.3	1665	60.3
	No	479	37.6	330	38.4	286	45.7	1096	39.7
The dangers of tobacco use?	Yes	841	66.3	558	65.5	357	57.4	1755	64.0
	No	427	33.7	294	34.5	265	42.6	987	36.0
When parents find out their kids have been drinking they may discuss it, take away privileges, add chores, take away cell phones, use of the car, etc. In the past 12 months, if your parents found out you were drinking, how did they usually respond? (Select one option).	I didn't drink.	1106	86.3	681	78.5	396	62.8	2185	78.6
	I drank but I was not caught.	37	2.9	60	6.9	105	16.6	203	7.3
	I was caught but there were no consequences.	12	0.9	17	2.0	61	9.7	89	3.2
	There were minor consequences.	12	0.9	31	3.6	31	4.9	75	2.7
	There were major consequences.	114	8.9	78	9.0	38	6.0	231	8.3
If the police caught you drinking, which of the following would most likely happen? (Select one option).	There would be no consequence.	79	6.2	98	11.2	75	11.9	253	9.1
	I would be given a warning and then let go.	96	7.5	73	8.4	59	9.3	228	8.2
	I would be taken home to my parents.	467	36.5	340	39.0	240	38.0	1047	37.6
	I would be arrested but would get no penalty.	156	12.2	94	10.8	57	9.0	306	11.0
	I would be arrested and the court would impose a penalty.	483	37.7	267	30.6	201	31.8	949	34.1
<i>For the following for questions, during the past 12 months, how many times has each of the following things happened?</i>									
You had problems at school or work because you had been drinking.	Never	1270	99.6	855	99.4	607	96.7	2733	98.9
	Once	3	0.2	2	0.2	12	1.9	17	0.6
	Twice	1	0.1	1	0.1	6	1.0	8	0.3
	Three or four times	0	0.0	2	0.2	2	0.3	3	0.1
	Five or more times	1	0.1	0	0.0	1	0.2	3	0.1

# Data Tables

**Table 11. (Continued) CSAP Questions**

Question	Response	Grade 8		Grade 10		Grade 12		Total	
		Number	Percent	Number	Percent	Number	Percent	Number	Percent
<i>For the following for questions, during the past 12 months, how many times has each of the following things happened?</i>									
You had problems with your friends because you had been drinking.	Never	1267	99.2	838	97.4	601	96.0	2705	97.9
	Once	5	0.4	17	2.0	16	2.6	39	1.4
	Twice	3	0.2	3	0.3	7	1.1	11	0.4
	Three or four times	3	0.2	2	0.2	1	0.2	6	0.2
	Five or more times	1	0.1	0	0.0	1	0.2	3	0.1
You had problems with someone you were dating because you had been drinking.	Never	1264	99.0	842	98.2	596	94.5	2701	97.7
	Once	11	0.9	11	1.3	21	3.3	44	1.6
	Twice	1	0.1	0	0.0	6	1.0	8	0.3
	Three or four times	1	0.1	4	0.5	7	1.1	11	0.4
	Five or more times	0	0.0	0	0.0	1	0.2	0	0.0
You were hung over.	Never	1253	98.2	817	94.6	517	82.1	2587	93.4
	Once	13	1.0	32	3.7	42	6.7	86	3.1
	Twice	5	0.4	6	0.7	28	4.4	39	1.4
	Three or four times	4	0.3	8	0.9	20	3.2	33	1.2
	Five or more times	1	0.1	1	0.1	23	3.7	25	0.9
You were sick to your stomach or threw up after drinking.	Never	1265	98.9	837	97.0	539	85.7	2641	95.3
	Once	6	0.5	17	2.0	46	7.3	69	2.5
	Twice	4	0.3	7	0.8	23	3.7	33	1.2
	Three or four times	1	0.1	1	0.1	12	1.9	14	0.5
	Five or more times	3	0.2	1	0.1	9	1.4	11	0.4
You got into a physical fight because you had been drinking.	Never	1263	99.4	853	99.4	615	97.5	2732	99.0
	Once	4	0.3	4	0.5	8	1.3	17	0.6
	Twice	3	0.2	0	0.0	3	0.5	6	0.2
	Three or four times	1	0.1	1	0.1	1	0.2	3	0.1
	Five or more times	0	0.0	0	0.0	4	0.6	3	0.1
You were drunk at school or work.	Never	1269	99.7	851	99.1	608	97.1	2728	98.9
	Once	3	0.2	5	0.6	10	1.6	19	0.7
	Twice	0	0.0	1	0.1	3	0.5	3	0.1
	Three or four times	1	0.1	2	0.2	2	0.3	6	0.2
	Five or more times	0	0.0	0	0.0	3	0.5	3	0.1



# DFC and Youth Perception Tables

**Table 12. Drug Free Communities Report \***

Outcomes	Definition	Grade 8		Grade 10		Grade 12		Total		Male		Female	
		Per.	Num.	Per.	Num.	Per.	Num.	Per.	Num.	Per.	Num.	Per.	Num.
<b>How do you feel about someone your age having one or two drinks of an alcoholic beverage nearly every day? †</b>	Neither Approve nor Disapprove	14.5	229	20.5	224	25.3	202	18.9	656	21.3	363	16.5	290
	Somewhat Disapprove	8.7	137	15.3	168	17.8	142	12.9	448	12.9	220	12.8	225
	Strongly Disapprove	66.8	1053	51.5	564	43.3	345	56.6	1963	52.6	895	60.4	1061
	Don't know or can't say	10.0	158	12.7	139	13.6	108	11.7	406	13.2	225	10.3	181
<b>How much do you think people risk harming themselves (physically or in other ways) if they: (Moderate risk or Great Risk)</b>	take one or two drinks of an alcoholic beverage (beer, wine, liquor) nearly every day?	69.1	1413	67.5	959	65.9	694	67.9	3066	64.4	1472	71.1	1584
	have five or more drinks of an alcoholic beverage once or twice a week?	71.9	1402	71.1	956	63.4	692	69.7	3050	65.7	1469	73.6	1571
	smoke 1 or more packs of cigarettes per day.	73.2	1412	72.2	963	72.1	696	72.6	3071	69.6	1478	75.5	1583
	smoke marijuana once or twice a week?	63.0	1410	48.2	961	32.7	692	51.5	3063	48.2	1475	54.7	1578
	use prescription pain relievers that are not prescribed to them?	73.1	1399	71.2	955	67.9	692	71.3	3046	68.3	1467	74.2	1569
<b>How wrong do your parents feel it would be for YOU to: (Wrong or Very Wrong)</b>	have one or two drinks of an alcoholic beverage nearly every day?	99.3	1275	98.5	860	96.0	627	98.3	2762	98.2	1319	98.4	1436
	smoke cigarettes	99.2	1274	98.3	861	94.4	629	97.8	2764	98.3	1321	97.4	1436
	smoke marijuana	97.6	1270	93.4	860	83.7	627	93.1	2757	93.1	1316	93.3	1434
	use prescription pain relievers that are not prescribed to them?	98.7	1284	97.9	861	96.5	630	97.9	2775	98.3	1323	97.6	1445
<b>How wrong do your friends feel it would be for you to: (Wrong or Very Wrong)</b>	have one or two drinks of an alcoholic beverage nearly every day?	95.5	1412	90.6	970	81.9	700	90.9	3082	90.4	1478	91.4	1595
	smoke cigarettes	96.7	1411	93.5	970	81.1	697	92.1	3078	91.8	1482	92.4	1587
	smoke marijuana	92.8	1399	75.6	960	55.4	697	78.9	3056	79.0	1471	78.8	1576
	use prescription pain relievers that are not prescribed to them?	96.2	1410	93.6	965	87.7	691	93.4	3066	94.1	1479	92.9	1578
<b>Past 30 day use of (at least one use in the Past 30 Days):</b>	Alcohol	4.9	1362	14.4	921	32.6	675	14.2	2958	11.8	1419	16.2	1528
	Cigarettes	0.9	1350	2.7	904	9.6	658	3.4	2912	3.5	1395	3.2	1508
	Marijuana	1.3	1349	6.0	900	19.5	652	6.9	2901	6.2	1390	7.4	1502
	Any Prescription Drug	0.8	1359	2.2	907	4.7	659	2.1	2925	1.6	1398	2.6	1518
<b>Average Age of Onset **</b>		Age	Num.	Age	Num.	Age	Num.	Age	Num.	Age	Num.	Age	Num.
	Alcohol	11.6	178	13.5	268	14.5	361	13.6	807	13.4	334	13.6	470
	Cigarettes	11.7	52	13.1	95	14.2	183	13.5	330	13.5	151	13.6	177
	Marijuana	12.0	47	13.9	147	14.6	282	14.2	476	14.0	214	14.3	259
	Prescription Pain Relievers	11.8	47	13.8	40	15.3	61	13.8	148	14.2	59	13.5	87

† For this question only, the numbers and percentages represent the students' answers for all response options.

\* The "Num." column represents the sample size (the number of youth who answered the question). The "Per." column represents the percentage of youth in the sample answering the question as specified.

\*\* For Average Age of Onset, "Num." represents the number of youth who reported any age of first use for the specified substance other than "Never Used."

# DFC and Youth Perception Tables

**Table 13. Youth Perceptions of Substance Use**

Now think about all the students in your grade at school. How many of them do you think:	Substance	Grade 8		Grade 10		Grade 12		Total	
		Number	Percent	Number	Percent	Number	Percent	Number	Percent
<b>a. smoke one or more cigarettes a day?</b>	None (0%)	912	53.6	388	33.3	260	30.8	1559	42.0
	Few (1-10%)	555	32.6	309	26.5	167	19.8	1032	27.8
	Some (11-30%)	138	8.1	255	21.9	151	17.9	546	14.7
	Half or less (31-50%)	58	3.4	120	10.3	134	15.9	312	8.4
	Half or more (51-70%)	22	1.3	63	5.4	72	8.5	156	4.2
	Most (71-90%)	10	0.6	17	1.5	39	4.6	67	1.8
	Almost All (91-100%)	7	0.4	14	1.2	21	2.5	41	1.1
<b>b. drank alcohol sometime in the past month?</b>	None (0%)	833	49.1	330	28.4	220	26.0	1381	37.3
	Few (1-10%)	507	29.9	134	11.5	57	6.7	696	18.8
	Some (11-30%)	212	12.5	186	16.0	91	10.8	489	13.2
	Half or less (31-50%)	83	4.9	200	17.2	119	14.1	404	10.9
	Half or more (51-70%)	41	2.4	156	13.4	134	15.9	330	8.9
	Most (71-90%)	15	0.9	127	10.9	156	18.5	296	8.0
	Almost All (91-100%)	7	0.4	29	2.5	68	8.0	104	2.8
<b>c. used marijuana sometime in the past month?</b>	None (0%)	914	54.1	346	29.7	229	27.1	1491	40.3
	Few (1-10%)	466	27.6	161	13.8	58	6.9	684	18.5
	Some (11-30%)	147	8.7	186	16.0	74	8.8	407	11.0
	Half or less (31-50%)	86	5.1	160	13.7	106	12.6	351	9.5
	Half or more (51-70%)	37	2.2	137	11.8	137	16.2	311	8.4
	Most (71-90%)	25	1.5	118	10.1	155	18.4	300	8.1
	Almost All (91-100%)	14	0.8	56	4.8	85	10.1	155	4.2
<b>d. used an illegal drug in the past month (not including marijuana)?</b>	None (0%)	1083	64.1	427	36.9	279	33.2	1789	48.5
	Few (1-10%)	456	27.0	296	25.6	197	23.4	952	25.8
	Some (11-30%)	68	4.0	193	16.7	118	14.0	380	10.3
	Half or less (31-50%)	46	2.7	109	9.4	100	11.9	254	6.9
	Half or more (51-70%)	15	0.9	61	5.3	63	7.5	140	3.8
	Most (71-90%)	14	0.8	54	4.7	54	6.4	122	3.3
	Almost All (91-100%)	8	0.5	17	1.5	30	3.6	55	1.5

### NATIONAL RESOURCES

United States Department of Health and Human Services (USDHHS)  
Substance Abuse and Mental Health Service Administration (SAMHSA)  
1 Choke Cherry Rd., Rm. 8-1054  
Rockville, Maryland 20857  
240-276-2000

[info@samhsa.hhs.org](mailto:info@samhsa.hhs.org)

[www.samhsa.gov](http://www.samhsa.gov)

(From this web-site, the programs and services provided by the Center for Substance Abuse Prevention, Center for Substance Abuse Treatment, and Center for Mental Health Services can be accessed)

Center for Substance Abuse Prevention (CSAP)

1 Choke Cherry Rd., Ste 4-1057  
Rockville, Maryland 20857  
240-276-2420

[info@samhsa.hhs.org](mailto:info@samhsa.hhs.org)

<http://prevention.samhsa.gov/>

CSAP's Centers for the Advancement of Prevention Technologies (all five CSAP Centers can be accessed through this web site)

<http://captus.samhsa.gov/home.cfm>

National Institutes of Health (NIH)  
6001 Executive Blvd., Rm. 5213  
Bethesda, Maryland 20892-9561  
301-443-1124

<http://www.nida.nih.gov/>

### STATE RESOURCES

SAFE, Inc.  
Chesterfield Youth Planning and Development  
9700 Krause Road  
Chesterfield, VA 23832  
804-267-3377

<http://www.chesterfieldsafe.org/>

### This Report Was Prepared by Bach Harrison L.L.C.

R. Steven Harrison, Ph.D.  
R. Paris Bach-Harrison, B.F.A.  
Taylor C. Bryant, B.A.

801-359-2064

<http://www.bach-harrison.com>