


# Assessment of mild illnesses in children




\*Symptoms of COVID-19: fevers/chills/rigor, cough, difficulty breathing, headache, myalgias, congestion/runny nose, nausea/vomiting/diarrhea, sore throat, loss of smell and/or taste

\*\*People who have tested positive for COVID-19 (laboratory confirmed) do not need to quarantine or get tested again for up to 3 months as long as they do not develop symptoms again

\*\*\* Fever-free means temperature less than 100.4 F without fever-reducing medications

--Monitor local flu epidemiology ([www.cdc.gov/flu/weekly](http://www.cdc.gov/flu/weekly)) and refer to IDSA guidance on diagnosis and treatment of flu ([www.idsociety.org/practice-guideline/influenza](http://www.idsociety.org/practice-guideline/influenza)). Coinfection with SARS-CoV-2 and influenza is possible. Data on co-infections with COVID-19 and other respiratory infections such as Group A strep is not available at this time

# Assessment of children with exposure to COVID-19



\*Symptoms of COVID-19: fevers/chills/rigor, cough, difficulty breathing, headache, myalgias, congestion/runny nose, nausea/vomiting/diarrhea, sore throat, loss of smell and/or taste

\*\*People who have tested positive for COVID-19 (laboratory confirmed) do not need to quarantine or get tested again for up to 3 months as long as they do not develop symptoms again

\*\*\*Fever-free means temperature less than 100.4 F without fever-reducing medications