

## Additional References **Lithium and Suicide Risk**

Baldessarini RJ, Tondo L. Lithium and suicidal risk. *Bipolar Disord.* 2008; 10: 114–115.

Baldessarini RJ, Tondo L. Meta-analytic comparison of antisuicidal effects of lithium versus anticonvulsants. *Pharmacopsychiatry.* 2009; 42: 72–75.

Baldessarini RJ, Tondo L, Davis P, Pompili M, Goodwin FK, Hennen J. Decreased risk of suicides and attempts during long-term lithium treatment: a meta-analytic review. *Bipolar Disord.* 2006; 8: 625–639.

Guzzetta F, Tondo L, Centorrino F, Baldessarini RJ. Lithium treatment reduces suicidal risk in recurrent major depressive disorder. *J Clin Psychiatry.* 2007; 68: 380–383.

Tondo L, Baldessarini RJ. Reduction of suicidal behavior in bipolar disorder patients during long-term treatment with lithium. Chapt 26 in Koslow SH, Ruiz P, Nemeroff CB (editors): *A Concise Guide to Understanding Suicide*. Cambridge, UK: Cambridge University Press, 2014, pp 217–228.

Tondo L, Baldessarini RJ. Suicide in bipolar disorder. Chapt 37 in *The Bipolar Book: History, Neurobiology, and Treatment*, edited by Yildiz A, Nemeroff C, and Ruiz P. New York: Oxford University Press, 2015, pp 509–528.

Tondo L, Hennen J, Baldessarini RJ. Reduced suicide risk with long-term lithium treatment in major affective illness: a meta-analysis. *Acta Psychiatr Scand.* 2001; 104: 163–172.

Tondo L, Vásquez GH, Baldessarini RJ. Prevention of suicidal behavior in bipolar disorder. *Bipolar Disord.* 2020.