



PRIME SOURCE

TRIAD **QC** SELECT™

Broad spectrum control of grassy and broadleaf weeds



ACTIVE INGREDIENT

2,4-D
Quinclorac
Dicamba

% ACTIVE INGREDIENT

2,4-D 13.29%
Quinclorac 9.79%
Dicamba 1.67%

FORMULATION

Liquid

CHEMICAL FAMILY

Group 4 herbicides
2,4-D - Phenoxy
Quinclorac - Carboxylic acids
Dicamba - Benzoic acids

BRAND ALTERNATIVE

Quincept®

SIGNAL WORD

Warning

PACKAGING

12 x 1 qt, 2 x 2.5 gal

LABELED USES

Residential and non-residential turfgrass, including lawns or grounds around residential establishments

Athletic fields

Parks

Roadsides

Golf course turf – not for use on greens, tees and collars

Sod farms

IMPORTANT:

ALWAYS READ AND FOLLOW LABEL INSTRUCTIONS.



Triad QC Select™ provides post-emergent control of a comprehensive list of both grassy and broadleaf weeds.

- Combines 3 active ingredients to help eliminate hard to control weeds.
- Perfect late season clean up product for areas that didn't receive pre-emergent herbicide application in the spring.
- Can be applied as a spot application or broadcast spray.
- Controls newly germinated 1- to 2-leaf crabgrass, to 1-tiller crabgrass and when crabgrass has matured to 5-tillers or greater.
- Presence of Quinclorac contributes to control of grassy weeds, is absorbed by the foliage and roots.

ORNAMENTAL TURF AND SOD

Broadcast Treatment: Apply at a rate of 7 to 8 pints/A (112 to 128 fl oz/A) in a spray volume of 20 to 300 gal/A (0.5 to 6.9 gal/1,000 sq ft). Do not exceed 2 broadcast applications per year, excluding spot treatments. For sod the minimum retreatment interval is 21 days.

Spot Treatment: Apply at a rate of 2.6 to 2.9 fl oz per 1,000 sq ft in a spray volume of 1 gal. Do not apply greater than 16 pints of this product per acre per year.

The minimum retreatment interval is 21 days.

TRIAD SELECT™

Broad spectrum control of grassy and broadleaf weeds

NON-TURF AREAS

Control of annual and perennial plants

Broadcast Treatment: Apply at a rate of 7 to 8 pints/A (112 to 128 fl oz/A) in a spray volume of 20 to 300 gal/A (0.5 to 6.9 gal/1,000 sq ft).

Do not exceed 2 broadcast applications per year, excluding spot treatments, with a minimum retreatment interval of 30 days.

Spot Treatment: Apply at a rate of 2.6 to 2.9 fl oz per 1,000 sq ft in a spray volume of 1 gal. Do not apply greater than 16 pints of this product per acre per year.

USE RESTRICTIONS

- Do not use on golf course greens, tees, and collars.
- Do not use on lawns with desirable clovers or legumes or on ornamentals.
- Not for use on sod farms in Arizona. For use in New York by spot treatment only (spray individual weeds only; adjust the sprayer to coarse spray to minimize wind drift, apply to the center of the weeds and spray to lightly cover).

Read label for full application and rate information.

Weeds controlled include:

Barnyard grass	Dock	Morning glory	Virginia buttonweed
Black medic	Dogbane	Nettle	Virginia creeper
Boxelder	Elderberry	Oxalis	White clover
Broadleaf dock	Fall dandelion	Plantain	Wild carrot
Burweed	Fiddleneck	Russian pigweed	Wild vetch
Carpetweed	Field bindweed	St. Johnswort	Willow
Chickweed	Ground ivy	Shepherds purse	Woodsorrel
Clover	Hedge bindweed	Smartweed	Yarrow
Crabgrass, large and smooth	Henbit	Spurge	Yellow rocket
Dandelion	Horsetail	Thistle	
	Knotweed	Trailing crown vetch	

Refer to label for full list of broadleaf and grassy weeds controlled.