

DPI GLOBAL PRODUCT USAGE SHEET

Micro-Aid® Feed Grade Concentrate

Swine	Ounces (oz)/Ton	Grams (g)/Metric Tonne
Starter	4 oz	125 g
Grower/Finisher	2 oz	62.5 g
Sows – Gestation	2 oz	62.5 g
Sows – Lactation	4 oz	125 g

Poultry	Ounces (oz)/Ton	Grams (g)/Metric Tonne
Layers	2 to 4 oz	62.5 to 125 g
Broilers	2 to 4 oz	62.5 to 125 g
Turkeys	4 oz	125 g

Beef	Body Weight, mg/kg or mg/lb	Daily, mg/head/day
Beef Cattle	2.2 mg/kg or 1 mg/lb	500 to 1,000 mg

Dairy	Daily, mg/head/day
Calves	500 to 1,000 mg
Cows	1,000 to 2,000 mg
Heifers	1,000 to 2,000 mg

Horses	Daily, mg/head/day
Horses	750 mg

Shrimp	Pounds (lbs)/Ton	Grams (g)/Metric Tonne
Shrimp	1 to 2 lb	500 to 1,000 g

Companion/Exotic	Ounces (oz)/Ton	Grams (g)/Metric Tonne
Cats	4 oz	125 g
Deer	4 oz	125 g
Dogs	4 oz	125 g
Goats	4 oz	125 g
Rabbits	4 oz	125 g

Micro-Aid® Liquid

Water/Lagoon Treatment

Initial rate= 10 gallons/Million Gallons
Weekly rate = 4 gallons/Million Gallons

Pit Treatment

4 oz/100 Cubic Feet Sludge

Cocci-Guard®

For use in making finished broiler feeds at the rate of 500 to 1,000 grams per metric tonne of starter ration (0 to 21 days) and 500 grams per metric tonne of grower/finisher ration (22 days to market)

ENDURANCE D.R.O.

Feed one 1 oz/head/day (1 scoop) on top of normal, grain-based ration

Odor-Bloc® Equine Stall Odor Eliminator

Sprinkle 1 pound per 100 square feet each time stall is cleaned and bedding is replaced. For extended use, periodically sprinkle a small amount on top of bedding to keep odors away longer.

Odor-Bloc® Cat Litter Deodorizer

Sprinkle 1/3 to 1/2 cup on top of fresh, clean litter. For extended use or multiple cat families, periodically sprinkle a small amount on top of litter to keep odors away longer.

SSO®

Apply 65 oz/acre at planting and again as a foliar treatment