

Ken Caryl Dentistry  
7631 Shaffer Pkwy Suite B  
Littleton, CO 80127

Phone Number: (303) 973-5280  
Email: [info@kencaryldentistry.com](mailto:info@kencaryldentistry.com)  
Website: [www.kencaryldentistry.com](http://www.kencaryldentistry.com)

## **AFTER DENTAL IMPLANT SURGERY** Surgical Instructions

Do not disturb the wound. Avoid rinsing, spitting, or touching the wound on the day of surgery. There will be a metal healing abutment protruding through the gingival (gum) tissue.

### **Bleeding**

Some bleeding or redness in the saliva is normal for 24 hours. Excessive bleeding (your mouth fills up rapidly with blood) can be controlled by biting on a gauze pad placed directly on the bleeding wound for 30 minutes. If bleeding continues please call for further instructions.

### **Swelling**

Swelling is a normal occurrence after surgery. To minimize swelling, apply an ice bag, or a plastic bag, or towel filled with ice on the cheek in the area of surgery. Apply the ice continuously, as much as possible, for the first 36 hours.

### **Diet**

Drink plenty of fluids. Avoid hot liquids or food. Soft food and liquids should be eaten on the day of surgery. Return to a normal diet as soon as possible unless otherwise directed.

### **Pain**

You should begin taking pain medication as soon as you feel the local anesthetic wearing off. For moderate pain, 1 or 2 Tylenol or Extra Strength Tylenol may be taken every 3-4 hours. Ibuprofen (Advil or Motrin) may be taken instead of Tylenol. Ibuprofen, bought over the counter comes in 200 mg tablets: 2-3 tablets may be taken every 3-4 hours as needed for pain. For severe pain, the prescribed medication should be taken as directed, instead of over the counter medications. Do not take any of the above medication if you are allergic, or have been instructed by your doctor not to take it.

### **Antibiotics**

Be sure to take the prescribed antibiotics as directed to help prevent infection

### **Oral Hygiene**

Good oral hygiene is essential to good healing. The night of surgery, use the prescribed Peridex Oral Rinse before bed. The day after surgery, the Peridex should be used twice daily, after breakfast and before bed. Be sure to rinse for at least 30 seconds then spit it out. Warm salt water rinses (teaspoon of salt in a cup of warm water) should be used at least 4-5 times a day, as well, especially after meals. Brushing your teeth and the healing abutments is no problem. Be very gentle initially with brushing the surgical areas. Avoid brushing areas with sutures.